

Synthesis, spectroscopic characterization and antimicrobial evaluation of some (*E*)-*N*-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

P. Jayanthi¹, G. Jesu Retna Raj¹, M. Sekar^{1,*} and V. Thanikachalam²

¹PG & Research Department of Chemistry, Government Arts College,
C. Mutlur, Chidambaram - 608102, India

²Department of Chemistry, Annamalai University, Annamalainagar - 608002, Tamil Nadu, India

*E-mail address: drmschemgac@gmail.com

ABSTRACT

In the present study, six numbers of Schiff bases (1-6) have been synthesized by the condensation of 4-fluorobenzenesulfonamide and substituted aromatic aldehyde. The purities of these Schiff bases have been checked by their physical constants, IR, ¹H NMR and ¹³CNMR spectral data. The antimicrobial activities of these Schiff bases have been evaluated using Bauer-Kirby method.

Keywords: 4-fluorobenzenesulfonamide, IR spectra, NMR spectra, Bauer-Kirby, antimicrobial activities

1. INTRODUCTION

Schiff base, an organic compound having general formula R-C=N-R' where R and R' are aryl, alkyl or cycloalkyl or heterocyclic groups formed by the condensation of an amine and a carbonyl group, is a potential inhibitor. The greatest advantage of many Schiff base compounds is that they can be conveniently and easily synthesized from relatively cheap material. Schiff base compounds due to the presence of the -C=N- group, electronegative nitrogen, sulfur and/or oxygen atoms in the molecule, have been reported to be effective

inhibitors for the corrosion of iron and steel in acidic and alkaline media by several authors^{1,12}.

Schiff bases exhibit excellent characteristics and structural similarities with natural biological substances, relatively simple preparation procedures and the synthetic flexibility that enables design of suitable structural scaffolds^{13,14}.

The Schiff bases are also used as versatile components in nucleophilic addition with organometallic reagents¹⁵ and in cycloaddition reactions^{16,17}. Schiff bases have wide applications in food industry, dye industry, analytical chemistry, catalysis, fungicidal, agrochemical and biological activities¹⁸ with the increasing incidence of deep mycosis, there has been increasing emphasis on the screening of new and more effective antimicrobial drugs with low toxicity¹⁹.

Schiff bases are associated with antibacterial, antifungal and anti tubercular activities have diverse biological activities²⁰⁻²⁴. The Schiff bases engrossed much attention as they demonstrated antimicrobial²⁵, anticancer²⁶, anticonvulsant²⁷, diuretic²⁸, herbicidal²⁹, anti-inflammatory³⁰, antitumor³¹ and anti HIV³² activities.


In the present investigation, the synthesis of some new Schiff bases from 4-fluorobenzenesulfonamide and substituted aromatic aldehyde. The synthesis of Schiff bases were characterized by IR, ¹H NMR and ¹³C NMR. The Schiff bases were also screened for their antimicrobial activities of the prepared compounds were assessed against Gram-positive bacteria, Gram-negative bacteria and fungi.

2. MATERIALS AND METHODS

All the chemicals involved in the present investigation, have been procured from Sigma-Aldrich and E-Merck chemical companies. Melting points of all Schiff bases have been determined in open glass capillaries on SUNTEX melting point apparatus and are uncorrected. Infrared spectra (KBr, 4000–400 cm⁻¹) have been recorded on Avatar-330 FT-IR spectrophotometer. The NMR spectra of all synthesized compounds have been recorded on Bruker 400 MHz spectrometer operating at 400 MHz for recording ¹H spectra and 100 MHz for ¹³C spectra in CDCl₃ solvent using TMS as internal standard.

General procedure for synthesis of

(E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides



Scheme 1. Synthesis of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

The Schiff bases were obtained by 0.1 M solution of 4-fluorobenzenesulfonamide was added to 0.1M solution of substituted aromatic benzaldehyde in ethanol, the addition reaction mixture is heated under reflux for 5-6 hours at 70 °C. After completion of reaction the precipitate are formed. The product filtered after cooling and purified with ethanol. The purity of product was checked by MP and TLC. The synthesized compounds were characterized by their Physical constants, IR, ¹H NMR and ¹³C NMR spectral data. The analytical and spectral data of synthesized Schiff bases are given below.

Spectral data of (E)-N-(4-fluorobenzylidene)-4-fluorobenzenesulfonamide (1)

IR($\nu_{\text{cm}^{-1}}$) = 1589.34 (C=N); ¹H-NMR (400 MHz, DMSO-d₆), ¹H(-NMR(δ ppm) = 8.420 (1H, s, CH=N); 7.090-7.933(12H, m, Ar-H); ¹³C-NMR (100 MHz, DMSO-d₆), ¹³C-NMR(δ ppm) = 175.767(C=N), 147.825(C₁) 130.745(C₂ & C₆), 116.015.(C₃ & C₅), 161.924 (C₄-F), 158.552(C₁₆-F), 130.774(C₁₃), 132.449(C₁₄ & C₁₈), 115.017(C₁₅), 115.869(C₁₇), M.F. C₁₃H₉F₂NO₂S, M.W. 281.0; m.p. 61–62 °C.

Spectral data of (E)-N-(4-chlorobenzylidene)-4-fluorobenzenesulfonamide (2)

IR($\nu_{\text{cm}^{-1}}$) = 1589.34 (C=N); ¹H-NMR (400 MHz, DMSO-d₆), ¹H(-NMR(δ ppm) = 8.422 (1H, s, CH=N); 7.187-7.975(8H, m, Ar-H); ¹³C-NMR (100 MHz, DMSO-d₆), ¹³C-NMR(δ ppm) = 175.653(C=N), 162.952.(C-F), 147.842(C₁), 132.496(C₂ & C₆), 115.912 (C₃), 115.907(C₅), 130.762(C₁₃), 130.777(C₁₄&C₁₈), 132.489(C₁₅&C₁₇), 147.818(C₁₆), (M.F.C₁₃H₉ClFNO₂S; M.W. 297.73 m.p. 59–60 °C.

Spectral data of (E)-N-(4-bromobenzylidene)-4-fluorobenzenesulfonamide (3)

IR($\nu_{\text{cm}^{-1}}$) = 1589.34 (C=N); ¹H-NMR (400 MHz, DMSO-d₆), ¹H(-NMR(δ ppm) = 8.391 (1H, s, CH=N); 7.092-7.780 (8H, m, Ar-H); ¹³C-NMR (100 MHz, DMSO-d₆), ¹³C-NMR(δ ppm) = 176.178(C=N), 162.100(C-F), 134.0847(C₁), 133.024(C₂ & C₆), 119.587(C₃ & C₅), 133.024(C₁₅), & C₁₇, 132.024(C₁₃), 124.33(C₁₄) & C₁₈, 122.508(C₁₆), M.F. C₁₃H₉BrFNO₂S, M.W. 342.18; m.p. 57–58 °C.

Spectral data of (E)-N-(4-methylbenzylidene)-4-fluorobenzenesulfonamide (4)

IR($\nu_{\text{cm}^{-1}}$) = 1589.34 (C=N); ¹H-NMR (400 MHz, DMSO-d₆), ¹H(-NMR(δ ppm) = 8.421 (1H, S, CH=N); 7.091-7.819 (8H, m, Ar-H); 2.48(3H, s, methyl-H), ¹³C-NMR (100 MHz, DMSO-d₆), ¹³C-NMR(δ ppm) = 175.411(C=N), 165.154(C-F), 21.697(CH₃), 148.247(C₁) 133.553 (C₂ & C₆), 128.789(C₃ & C₅), 129.527(C₁₃), 141.967(C₁₄&C₁₈), 128.789(C₁₅ & C₁₇), 148.262(C₁₆), M.F. C₁₄H₁₂FNO₂S; M.W.277.31; m.p. 55-56 °C.

Spectral data of (E)-N-(4-methoxybenzylidene)-4-fluorobenzenesulfonamide (5)

IR($\nu_{\text{cm}^{-1}}$) = 1589.34 (C=N); ¹H-NMR (400 MHz, DMSO-d₆), ¹H(-NMR(δ ppm) = 8.392(1H, S, CH=N); 7.001-7.8274(8H, m, Ar-H); 3.982(3H, s, methoxy H), ¹³C-NMR (100 MHz, DMSO-d₆), ¹³C-NMR(δ ppm) = 175.822(C=N), 165.412(C-F), 55.489(OCH₃), 148.312 (C₁) 129.149(C₂ & C₆), 115.841,.(C₃ & C₅), 122.269(C₁₃), 130.428(C₁₄ & C₁₈), 114.217(C₁₅ & C₁₇), 161.745 (C₁₆), M.F. C₁₄H₁₂FNO₃S, M.W.293.31; m.p. 57-58 °C.

Spectral data of (E)-N-(4-nitrobenzylidene)-4-fluorobenzenesulfonamide (6)

IR(vcm^{-1}) = 1589.34 (C=N); $^1\text{H-NMR}$ (400 MHz, DMSO- d_6), $^1\text{H-NMR}(\delta \text{ ppm})$ = 8.399 (1H, S, CH=N); 7.101-7.8271(8H, m, Ar-H); $^{13}\text{C-NMR}$ (100 MHz, DMSO- d_6), $^{13}\text{C-NMR}(\delta \text{ ppm})$ = 175.706(C=N), 165.359(C-F), 150.400(C-NO₂), 145.663(C₁), 132.315(C₂ & C₆), 121.246 (C₃ & C₅), 134.242(C₁₃), 130.588(C₁₄& C₁₈), 127.170(C₁₅&C₁₇), M.F.C₁₃H₉FN₂O₄S, M.W. 308.28; m.p. 109–110 °C.

3. ANTIMICROBIAL ACTIVITIES

Antibacterial activity

(E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides(1-6) were tested for their antibacterial activity against two gram positive pathogenic strains *Bacillus subtilis*, *Staphylococcus aureus* and two gram negative strains *Escherichia coli* and *Pseudomonas aerogenosa*. The disc diffusion technique was followed using the Kirby–Bauer³³ method, at a concentration of 250 mg/mL with ciprofloxacin taken as the standard drug. The antibacterial screening effect of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides is shown in Fig-1 (Plates1–4). The measured zone of inhibitions is shown in Table-1 and the clustered column chart in Fig-2.

Results showed that most of the compounds possess potent to moderate activity as compared to the reference drug *ciprofloxacin*. Schiff base analogues (1-6) showed potent activity against Gram positive and significant activity against Gram negative bacterial strains. Compound (2) & (3) having 4-chloro and 4-bromo phenyl group in Schiff base exhibited potent activity against *Escherichia coli*, *Pseudomonas aeruginosa* and *Staphylococcus aureus*. while significant activity against *Bacillus subtilis*.

Plate – 1



Plate – 2



Plate – 3





Plate – 4




Fig. 1. Petri-plates (1-4) for antibacterial activities of (*E*)-*N*-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

When bromo or chloro group was replaced by fluoro or nitro group (1) & (6) the activity was enhanced for *Bacillus subtilis*. while it was diminished for *E. coli* and *Staphylococcus aureus*. Compound (5) showed significant activity against only *Pseudomonas aeruginosa* with 25 mm, and also, moderate activity was noticed against other strains. Compounds (2), (3) and (4) revealed better activity in comparison to other compounds used in study, indicating that methyl and halogen substitution at 4-position of Schiff base nucleus showed better activity as compared to standard.

Table 1. Zone of Inhibition (mm) values of antibacterial activities of (*E*)-*N*-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

S. No.	Substituents	Zone of Inhibition (mm)			
		Gram positive Bacteria		Gram negative Bacteria	
		<i>B. subtilis</i>	<i>S. aureus</i>	<i>E. coli</i>	<i>Pseudomonas</i>
1	4-F	31	-	16	23
2	4-Cl	20	28	19	24
3	4-Br	14	25	29	25

4	4-CH ₃	22	12	24	25
5	4-OCH ₃	8	15	21	24
6	4-NO ₂	31	16	13	25
Standard	Ciprofloxacin	33	21	29	31
Control		-	-	-	-




Fig. 2. Cluster Column for Antibacterial activity of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

Antifungal activity

The antifungal activities of all the synthesized compounds have been studied against *Tricodermaviridi*, *Aspergillusniger*, *Muccor species* and *Candida albicans*. The disc diffusion technique was followed using the Kirby–Bauer³³ method, at a concentration of 250 mg/mL with Miconazole taken as the standard drug. The antifungal activities of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides have been studied and are shown in Fig-3. Plates (5-8) and the zone of inhibition values is given in Table-2 and the Clustered column Chart given in Fig-4.

Thus the substituents place a vital role in imparting enhanced antifungal activity to the compounds. Compounds (1), (2) & (3) were found to have better antifungal activities than those of other compounds. The higher activity of (1), (2) & (3) were influenced by the

presence of electron withdrawing group such as halogen on phenyl ring. Compound (4) having methyl group on para position of phenyl ring showed increased activity (19mm) against *Candida albicans* and similarly the compound (5) having methoxy group on para position of phenyl ring showed (19 mm) against *Aspergillus niger*. Among these compounds, compound (3) has strong selectivity towards all the fungal strains, but compound (2) has strong activity (26mm) against *Tricoderma viridi*.

Plate – 5




Plate – 6




Plate – 7



Plate – 8




Fig. 3. Petri-plates for antifungal activities of (*E*)-*N*-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

Table 2 Zone of Inhibition (mm) values of antifungal activities of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

S. NO.	Substituents	Zone of Inhibition (mm)			
		<i>Tricoderma viridi</i>	<i>Aspergillus niger</i>	<i>Mucor species</i>	<i>Candida albicans</i>
1	4-F	10	23	18	11
2	4-Cl	26	20	18	20
3	4-Br	21	22	22	21
4	4-CH ₃	14	18	11	19
5	4-OCH ₃	16	19	14	15
6	4-NO ₂	18	18	14	15
Standard	Miconazole	28	24	22	29
Control	DMSO	-	-	-	-




Fig. 4. Cluster Column for Antifungal activity of (E)-N-(4-substitutedbenzylidene)-4-fluorobenzenesulfonamides

4. CONCLUSION

A six new schiff bases have been synthesized by condensation method. These schiff bases have been characterized by their physical constants, IR, ¹HNMR & ¹³CNMR spectral data. The antibacterial activity of all synthesized schiff bases have been studied using Bauer-Kirby method. Most of the synthesized Schiff bases have shown better activity against Gram positive, Gram negative bacterial and Fungal Species compared to standard drugs Ciprofloxacin and Miconazole.

Acknowledgment

The authors thank DST NMR facility, Department of Chemistry, Annamalai University, Annamalai Nagar - 608 002, for recording NMR spectra of all compounds

References

- [1] M. Behpour, S.M. Ghoreishi, A. Gandomi-Niasar, N. Soltani and M. Salavati-Niasari (2009). *J. Mater. Sci.* 44, 2444.
- [2] A. Asan, S. Soylu, T. Kiyak and F. Yıldırım (2006). *Corros. Sci.* 48, 3933.
- [3] M. Behpour, S.M. Ghoreishi, N. Soltani, M. Salavati-Niasari, M. Hamadani and A. Gandomi (2008). *Corros. Sci.* 50(8), 2172.
- [4] A.M. Abdel-Gaber, M.S. Masoud, E.A. Khalil and E.E. Shehata (2009). *Corros. Sci.* 51, 3021.
- [5] S. Chitra, K. Parameswari and A. Selvaraj (2010). *Int. J. Electrochem. Sci.* 5, 1675.
- [6] R. Solmaz (2010). *Corros. Sci.* 52, 3321.
- [7] R. Solmaz, Ece Altunbaş and G. Kardas (2011). *Mater. Chem. Phys.* 125, 796.
- [8] M.G. Hosseini, M. Ehteshamzadeh and T. Shahrabi (2007). *Electrochim. Acta* 52, 3680.
- [9] R. Álvarez-Bustamante, G. Negrón-Silva, M. Abreu-Quijano, H. Herrera-Hernández, M. Romero-Romo, A. Cuán and M. Palomar-Pardavé (2009). *Electrochim. Acta* 54, 5393.
- [10] N.A. Negm and M.F. Zaki (2008). *Colloid Surf.* 322(A), 97.
- [11] M. Ehteshamzadeh, A.H. Jafari, N. Esmaeel and M.G. Hosseini (2009). *Mater. Chem. Phys.* 113, 986.
- [12] H. Shokry, M. Yuasa, I. Sekine, R.M. Issa, H.Y. El-Baradie and G.K. Gomma (1998). *Corros. Sci.*, 40, 2173.
- [13] S. Patai Ed., *The Chemistry of the Carbon-Nitrogen Double Bond*, J. Wiley & Sons, 1970, London.
- [14] E. Jungreis and S. Thabet. Marcell Dekker, 1969, New York.

- [15] V. V. Kuznetsov, A. R. Palma, A. E. Aliev, A. V. Varlamov and N. S. Prostavkov, *Zh. Org. Khim.* (1991), 127, 1579.
- [16] A. Taggi, A. M. EHafez, H. Wack, B. Young, D. Ferrari and T. Lectka. *J. Am. Chem. Soc.* (2002), 124, 6626.
- [17] O. Tsuge and R. Kanemasa, *Adv. Heterocycl. Chem.* (1989) 45231.
- [18] N. L. Owen and M. V. S. Sultanbawa. *J. Chem. Soc.* (1949) 3098.
- [19] M. J. Gemi, C. Biles, B. J. Keiser, S. M. Poppe, S. M. Swaney, W. G. Tarapley, D. L. Romeso and Y. Yage. *J. Med. Chem.* (2000), 43(5), 1034.
- [20] W. A. Al-Masoudi, H. Tooama, J. Hammed, *Basrah J. Vet. Res.* (2014) 7, 33,
- [21] W. O. Foye, Principles of Medicinal Chemistry, 3rd edition, Varghese Publishing House, Bombay, (1989) 728.
- [22] Z. Y. Guo, R. Xing, S. Liu, Z. Zhong, X. Ji, L. Wang and P. C. Li. *Carbohydrate Res.* (2007), 342(10), 1329.
- [23] S. J. Wadher, M. P. Puranik, N. A. Karande and P. G. Yeole. *Int. J. Pharm. Tech. Res.* (2009), 1(1), 22.
- [24] C. Spinu, M. Pleniceanu and C. Tigae, *Turk. J. Chem.* (2008) 32, 487.
- [25] H. Temel and H. Hosgoren. *Trans. Met. Chem.* (2002) 27(6), 609.
- [26] A. Yildiz, B. Kiraz and Dülger. *J. Serb. Chem. Soc.* (2007) 72, 215.
- [27] R. Gudipati, R. N. R. Anreddy and S. Manda. *Saudi Pharm. J.* (2011). 19, 153.
- [28] M. Verma, S. N. Pandeya, K. N. Singh and J. P. Stables. *Acta Pharm.* (2004), 54, 49.
- [29] S. Ghosh, S. Malik, B. Jain and S. A. Iqbal. *J. Saudi Chem. Soc.* (2012), 16, 137.
- [30] W. Meiyi, L. Zhengming and L. Yonghong. *Chin. J. Org. Chem.* (2010), 30, 877.
- [31] A. Pandey, R. Rajavel, S. Chandraker and D. Dash. *E-J. Chem.* (2012), 9, 2524
- [32] F. Shabani, L. A. Saghatforoush and S. Ghammamy. *Bull. Chem. Soc. Ethiop.* (2010), 24, 193.
- [33] A. W. Bauer, W. M. M. Kirby, J. C. Sherris and M. Truck. *Am. J. Clin. Pathol.* (1996), 45, 493.

(Received 18 August 2017; accepted 07 September 2017)