

Trace metals analysis of soil and edible plant leaves from abandoned municipal waste dumpsite in Owerri, Imo state, Nigeria

Ibe Francis Chizoruo^{1,*}, Beniah Obinna Isiuku² and Enyoh Christian Ebere³

^{1,3}Group Research in Analytical Chemistry, Environment and Climate Change (GRACE&CC), Department of Chemistry, Imo State University, Owerri, Imo State, P.M.B 2000, Nigeria

²Department of Chemistry, Imo State University, Owerri, Imo State, P.M.B 2000, Nigeria

¹⁻³Corresponding Author: francispavo@yahoo.com*, obinnabisiuku@yahoo.com, cenyoh@gmail.com

ABSTRACT

Municipal refuse may increase heavy metal concentration in soil, even at low levels, and their resulting long-term cumulative health effects are among the leading health concerns all over the world. In this study, the concentration of heavy metals in soils and edible plant leaves grown in an abandoned dumpsite along Akachi road in Owerri municipal were investigated. The soil samples were collected at each plot using a soil auger at the depth of 0-10 cm. Leaves dominant edible plant species were selected and collected from each sample plot. The samples were dried in an oven with forced air at 40 °C, milled to fine powder then digested with 10 ml concentrated HNO₃ and 5 ml concentrated HClO₄ and were analyzed for Cr, Cu, Fe, Mn, Al, and Zn using H183200 MultiParameter Bench Photometer. Result showed metals in the ranges in soils; Cr: 150-280 >Fe: 116.50-203 >Cu: 12.4-18.8 >Mn: 0-20 >Al: 0.08-0.16 >Zn: 0-1.4 mg kg⁻¹ Dw and levels of metals in the edible plant leaves are in the order of: Zn>Fe>Cu>Al>Mn>Cr. Zn in particular was higher than FAO/WHO recommended limit. Application of Pollution Load Index and Ecological risk models showed the area is unpolluted and safe for use. Daily Metal Intake estimates showed that zinc is mostly consumed from the plant species. The trends in Transfer Factor for the heavy metal in vegetable samples studied were in order: Zn>Al>Cu>Mn>Fe>Cr. Therefore, abandoned solid waste dumpsites contained significant concentrations of heavy metals which are later absorbed and accumulated by plants growing it.

Keyword: Accumulation, Metal intake, Pollution indices, Risk, Zinc

1. INTRODUCTION

There have been increasing attentions in the last few decades in both developing and developed countries on the environmental problem of soil pollution associated with heavy metals (Wu *et al.*, 2014). This is assisted by the unorganized increasing urban expansion, industrial developments coupled with inadequate waste management, which causes significant alterations in the physical environment and increases accumulation of municipal waste. Soil which enables growing of plant (Enyoh *et al.*, 2017) and human activities is mostly affected. Urban areas are known for high level of industrial activities which generate more pollutants (Tanee and Eshalomi-Mario, 2015). An example of such area is Owerri municipal, which serve as a connecting route to Port Harcourt, River State, Ontisha, Anambra State and Aba, Abia State. For this reason, the city has high exponential increase in human population and boost for commercial activities, which persistently creates significant volumes of waste on daily basis. The city lacks proper solid waste regulations and proper disposal facilities, for harmful waste. The poor management of waste or effluents could create a number of adverse environmental impacts as waste may be toxic or radioactive (Onibokun and Kumuyi, 1996; Wong *et al.*, 2003; UNDP, 2006; Kimani, 2007; Verla *et al.*, 2017a). Furthermore, explosion may occur as waste produce methane gas when organic materials are decomposing and also produce leachates, which pollute and ruins the aesthetic quality of the soil (Cointreau Levine, 1997; Oyelola *et al.*, 2009). Studies have shown municipal refuse may increase heavy metal concentration in soil and underground water (Albores *et al.*, 2000; Okoronkwo *et al.*, 2005; 2006), even at low levels, and their resulting long-term cumulative health effects are among the leading health concerns all over the world (Oluyemi *et al.*, 2008, Njoku and Ibe, 2009). However, abandoned waste dumpsites in southeastern Nigeria and other parts of the country have been used extensively as fertile grounds for cultivating vegetables and plant based foodstuff, though research has indicated that the vegetables are capable of accumulating high levels of heavy metals from contaminated and polluted soils (Garcia *et al.*, 1981; Xiong, 1998; Obasi *et al.*, 2013). In Owerri city, most abandoned landfill sites are currently used for crop cultivation, rearing livestock, as playground for children and as sites for unauthorized settlements. This practice is scientifically unacceptable in this era and as such, there is need for proper assessment of dumpsite waste soils to ensure environmental sustainability (Obasi *et al.*, 2013). Though numerous studies have been reported on heavy metals from dumpsites soils and crops grown on them (Amusan *et al.*, 2005; Okoronkwo *et al.*, 2005; Ololade *et al.*, 2007; Elaigwu *et al.*, 2007; Ndukwu *et al.*, 2008; Ebong *et al.*, 2008; Goorah *et al.*, 2009; Ayan *et al.*, 2010; Obute *et al.*, 2010; Charlse *et al.*, 2011; Magaji, 2012; Obasi *et al.*, 2012; Kanmani and Gandhimanti, 2013; Leah and Johny, 2014; Osei-Akoto *et al.*, 2016). However, the need for this research is due to paucity of data as well as published work regarding the fate of trace metal in this abandoned waste dumpsite. Since people now leave in this area and cultivates the land. This study will build on these works by attempting to address part of this problem by assessing the levels of heavy metals in the abandoned dumpsite located at Akachi road, Owerri municipal. This will involve determination of the concentration of these heavy metals in soil and edible plant leaves as well as subjecting them to index models for risk assessment. It is expected that results obtained from this study will widen our knowledge on the environmental risks associated with solid waste dumps in terms of heavy metal toxicity and the suitability of such site for plant cultivation. This will be of great benefit to the immediate community, the public and environmental health policy makers.

2. MATERIAL AND METHOD

2. 1. Study location

The study was conducted in Owerri municipal, which is one of the three Local Government areas that make up Owerri, the capital of Imo state in the Eastern part of Nigeria. According to 2006 population census of the Federal Republic of Nigeria, Owerri has a population of 401,873, (FRN, 2006) which must have increased tremendously over the past decade. Owerri Metropolis is located within latitudes 4°23'N - 7°15'N and longitudes 6°50' E and 7°25'E. The town is the capital of Imo State, having three big markets with a lot of commercial activities.

The location of the abandoned waste dump site is at Akachi road, Owerri, Imo state, Nigeria, as shown in fig. 1. This area was specifically chosen for the study because the waste dump site has been abandoned for over ten years. Some of the municipal wastes dumped in this site are now buried in the soil. There have been a lot of activities in this area in the recent years owing to the construction of a bypass across the abandoned waste dump site which opened up the area. The opening up of this area resulted in construction of both residential and commercial buildings. This is in addition to farming and growing of edible crops due to a river that passed through the area as shown in the legend (Fig. 1).




Figure 1. Map of Owerri showing sample location

2. 2. Sample collection and preparation

The sampling sites were spread objectively and the collections were done using a soil auger at 10 cm depth (top soil). At each site a “W” shaped line was drawn on a 2×2 m surface along which five samples were collected from each sample field area and mixed homogeneously to form one sample (Enyoh *et al*, 2017). The containers were washed with hot water and detergent and thoroughly rinsed in order to remove all contaminant.

The dominant plant species were collected using a knife from the abandoned site. Washed in distill water to remove soil/dust particles. They were put in black polyethene bags, tied and labeled with a masking tape and marker. Soil and plant samples were marked A to E to represent the same sample location. The soil and plant leaves were then transferred to the Imo State University Chemical laboratory for analysis. The plant leave samples were dried in an oven with forced air at 40 °C then they were milled to obtain fine powder and stored in a previously cleaned plastic bottle for heavy metals analysis. The samples were collected at five different points within the abandoned waste dump site at Akachi road. Coordinates of the sample collection points were referenced with GPS map 76 as presented (Table 1).

Table 1. Showing coordinates, elevation and description of sample area

SN	Sample site	Co-ordinate	Elevation (m)	Description of area
1	A	05°28'18.6" N 007°02'21.4" E	74	Along Akachi road, in a residential building
2	B	05°27'57.0" N 007°02'37.1" E	77	Along Akachi Road in a car wash
3	C	05°28'0.62" N 007°02'36.4" E	79	KenuJ 0 ² shopping mall
4	D	05°28'02.2" N 007°02'23.5" E	78	Njoku saw mil (a mini market)
5	E	05°28'21.8" N 007°02'27.3" E	67	Along Akachi road, at car wash

2. 3. Determination of Al, Cr, Cu, Fe, Mn and Zn

The soil and plant samples were digested with a mixture of nitric and perchloric acid. 2 g of powdered samples were accurately weighed into a beaker and then 10 ml of HNO₃ and 5 ml of HClO₄ was added, then it was allowed to stand for 24 hours. This was followed by filtration into a 100 ml volumetric flask and filtrate was made up to mark and used for analysis. The concentrations of metals were determined using Multiparameter Bench Photometer HI 83200 by HANNA Instruments (Duru *et al*, 2017).

2. 4. Analysis of potential ecological risk index:

The Potential Ecological Risk Index (RI) was assessed according to (Hakanson, 1980 and Aktaruzzaman *et. al.*, 2013) as the degree of heavy metal pollution in soil and according

to the toxicity of metals and the response of the environment (Aktaruzzaman *et. al.*, 2013). RI could evaluate ecological risk caused by toxic metals comprehensively. The calculating methods of RI are listed below

$$P_i = \frac{C_m}{C_o} \quad (1)$$

$$E_T = T_{ri} \times P_i \quad (2)$$

$$R_I = \sum ET_i \quad (3)$$

where, P_i is the single metal pollution index; C_m is the concentration of metal in the samples; C_o is the reference value for the metal; The reference concentrations were taken as the target value (mg/kg) of the Department of Petroleum Resources. The values are Fe = 38000, Zn = 140, Cu = 36, Cr = 100, and Mn = 850 (DPR, 2002; Verla *et. al.*, 2017b). E_T^i is the monomial potential ecological risk factor; T_{ri}^i is the metal toxic response factor according to Hakanson (1980). The values for each element are in the order Zn = 1, Cr = 2, Cu = 5. RI is the potential ecological risk caused by the overall contamination.

2. 5. Daily metals intakes estimate (DI_{metal})

The daily metals intakes were estimated by multiplying the concentration of metals in the edible plant leave with the recommended daily intake of at least 0.4 mg of vegetables (Sharma *et al.*, 2008).

$$DI_{metal} = C_{plant} \times 0.4 \quad (4)$$

C_{plant} is the concentration of metals in the plant samples, 0.4 is a factor recommended by experts (Sharma *et al.*, 2008).

2. 6. Transfer factors (TF) of the heavy metals from soils to the edible plant leaves

The transfer coefficient was calculated by dividing the concentration. The transfer coefficient was calculated by dividing the concentration of heavy metals in vegetables by the total heavy metal concentration in the soil (Kachenko and Singh, 2006).

$$TF = \frac{C_{plant}}{C_{soil}} \quad (5)$$

where, C_{plant} = metal concentration in plant tissue, mgkg⁻¹ fresh weight and C_{soil} = metal concentration in soil, mg kg⁻¹ dry weight (mg kg⁻¹ Dw)

3. RESULTS AND DISCUSSION

Heavy metal concentrations in the abandoned dumpsitesoil samples collected are presented in Table 2. The metal concentration ranges were: Cr: 150-280; Cu: 12.4-18.8; Fe: 116.50-203; Mn: 0-20; Al: 0.08-0.16; Zn: 0-1.4 mg kg⁻¹ Dw. The mean values were 264, 15.7, 169.3, 20, 0.12, 1.07 and standard deviations were 113.93, 2.62, 36.42, 0, 0.03, and 0.31 respectively.

Table 2. Metal concentrationin soil samples.

Metals (mg/kg)	Samples					Mean	SDV
	A	B	C	D	E		
Cr	210	230	450	150	280	264	113.93
Cu	17.3	16.3	18.8	13.7	12.4	15.7	2.62
Fe	203	202	116.50	152.00	173.00	169.3	36.42
Mn	ND	20	ND	20	20	20	0
Al	0.10	0.08	0.14	0.16	0.10	0.12	0.03
Zn	1.4	0.80	ND	ND	1.00	1.07	0.31

The concentrations of these elements obtained in this study were compared with values reported in literature. Chromium (Cr) concentration recorded in the present study (264 mg/kg) is highwhen compared to valuesrecorded in other studies such as 21.51 mg kg⁻¹ in the soil of Ado Ekiti, Southwestern Nigeria (Aruleba and Ajayi, 2012). Li *et al.* (2001) recorded 54.88 mg kg⁻¹ in urban soil of Hongkong. 22.27 mg/kg was observed in abandoned dumpsite in Umuahia, Abia State, Southeastern Nigeria (Okoronkwo et al, 2006). 4.55-33.46 mg/kg was recordedin the Region of DhakaAricha Highway, Savar, Bangladesh (Aktaruzzaman et al, 2013). In are related study carried out in China, 159.3 mg kg⁻¹ was recorded in urban soils and roadside dust (Shi *et al.*, 2008), while Jiang et al (2014) reported a mean value of 66.48 mg/kg in soils around coal gangue dump. The high level of chromium could be due to the nature of waste and persistent nature of the metal. The higher concentration values of chromium compounds could have been leached by rainwater and migrated through cracks in soil, asphalt roadways, and masonry walls, forming high-content chromium crystals on their surfaces (ATSDR, 2008). Chromium is carcinogenic resulting in cancer of respiratory organs in workers exposed to chromium containing dust (Langard, 1980).

The investigation found lower concentration of Cu compared to that in the soil of Gazipur (55-65) mg kg⁻¹ as reported by Habib *et al.* (2009) and soil around a coal gangue dump (22.72 mg/kg) reported by Jiang et al (2014). Though higher than mean values reported in soils of Bangladesh (9.92 mg/kg) as reported by Aktaruzzaman et al, (2013). In general, Cu

concentrations in the investigated soil were unpolluted since it is below the maximum acceptable limit of 70-80 mg/kg set by Federal Environmental Protection Agency, FEPA (1997).

Higher mean values of Fe were recorded in the present study when compared with the report of Osakwe and Okolie, (2015) who observed a mean value of 142.93 mgkg⁻¹. High concentrations of iron in soils relative to other metals have been reported in various studies (Dara, 1993; Ademoroti, 1996; Aluko and Oluwande, 2003) confirming that natural soils contain significant levels of iron. Furthermore, the value obtained in this study was far lower than value (12590.67 mg/kg) reported by Osei *et al.* (2016) in surface soils from abandoned waste sites in Ghana. Fe also was lower than the maximum acceptable limit of 400 mg/kg set by Federal Environmental Protection Agency, FEPA (1997). Lower level of Mn and Zn were obtained in the investigated soil when compared to mean values obtained by Chukwuocha *et al.*, (2015) who reported 150.7 mg/kg and 268.67 mg/kg for Mn and Zn respectively in top soils of Owerri, Imo state, Nigeria. The concentrations of metal found in soil of the present study were considerably significant and were moderately to heavily polluted for Cr, Cu, Mn and Fe according to soil quality guidelines for heavy metal in soil (Pekey, 2006). This observation suggests that dumping of waste materials might have influenced levels of these metals in the sites. Persons living or working in the vicinity of the sites may have been exposed through inhalation, ingestion, or skin contact with contaminated soils and dusts (ATSDR, 2008). On the other hand, Al and Zn in all the soil samples were unpolluted.

The degree of contamination is defined as the sum of all contamination factors. According to (Tomlinson *et al.*, 1980; Mathias and Stephen, 2016), the following terminologies are used to describe the degree of contamination; $P_i < 6$, low degree of contamination; $6 \leq P_i < 12$, moderate degree of contamination; $12 \leq P_i < 24$, considerable degree of contamination; $P_i \geq 24$, very high degree of contamination. In the present study, all sites were uncontaminated. The pollution load index is a potent tool in heavy metal pollution evaluation that provides a simple and comparative means for assessing the level of heavy metal pollution. The PLI represents the number of times by which the metal content in the soil exceeds the average natural background concentration, and gives a summative indication of the overall level of heavy metal toxicity in a particular sample.

Table 3. Individual pollution index or contamination factors for various metals studied.

Site	Individual Pollution Index						Degree	PLI	Comment
	Cr	Cu	Fe	Mn	Zn	$\sum P_i$			
A	0.7	0.48	0.0053	-	0.01	1.20	Low	0.02	No pollution
B	0.77	0.45	0.0053	0.02	0.0057	1.25	Low	0.002	No pollution
C	1.5	0.52	0.003	-	-	2.02	Low	0.24	No pollution
D	0.5	0.38	0.004	0.02	-	0.90	Low	0.02	No pollution
E	0.93	0.34	0.0045	0.02	0.007	1.30	Low	0.002	No pollution

*PLI = Pollution Load Index

The PLI value of > 1 is polluted, <1 indicates no pollution whereas values of $PLI = 1$ indicate heavy metal loads close to the background level (Cabarera *et al*, 1999; Mathias and Stephen, 2016). Therefore, the five sample points could be considered unpolluted for the metals studied.

Table 4. Monomial potential ecological risk factors and potential ecological risk caused by the overall contamination for various metals studied.

Site	Cr	Cu	Fe	Mn	Zn	$RI = \sum P_i$
A	1.4	2.4	-	-	0.01	3.81
B	1.54	2.25	-	-	0.0057	3.89
C	3	2.6	-	-	-	5.6
D	1	1.9	-	-	-	2.9
E	1.86	1.7	-	-	0.007	3.57

The following terminologies are used to describe the ecological risk factor: $E_{Ti} < 40$, low potential ecological risk; $40 \leq E_{Ti} < 80$, moderate potential ecological risk; $80 \leq E_{Ti} < 160$, considerable potential ecological risk; $160 \leq E_{Ti} < 320$, high potential ecological risk; and $E_{Ti} \geq 320$, very high ecological risk and the potential ecological risk index as given by Mathias and Stephen, (2016): $RI < 150$, low ecological risk; $150 \leq RI < 300$, moderate ecological risk; and $RI > 600$, very high ecological risk. As shown by table 4, all sample sites have low potential ecological risk, as well as low ecological risk. These suggest that the study area is safe for use.

Table 5. Metal concentration in plant samples

Metals (mg/kg)	<i>Carica Papaya</i> [1]	<i>Colocasia Esculenta</i>	<i>Psidium guajava</i>	<i>Prinus dulcis</i>	<i>Carica Papaya</i> [2]	Mean	SDV
Cr	0.015	0.07	0.085	0.1	0.15	0.08	0.05
Cu	3.0	3.55	4.25	4.00	4.20	3.80	0.53
Fe	16.40	1.05	3.25	3.85	10.20	6.95	6.28
Mn	0.5	ND	ND	1.50	1.50	1.17	0.57
Al	4.20	3	ND	5.00	1.05	3.32	1.72
Zn	225	75	75	100	150	125	63.74

*ND = Not Detected

Heavy metal concentrations in leafy vegetables of the edible plants grown on the abandoned dumpsite soil and their statistical analysis are presented in Table 5. Concentrations of heavy metal ranges were: Cr: 0.015-0.15; Cu: 3.0-4.25; Fe: 1.05-16.40; Mn: 0.5-3.1.50 mg kg⁻¹; Al: 1.05-5.00 and Zn: 75- 225 and order of heavy metals concentration in the vegetables samples were: Zn>Fe>Cu>Al>Mn>Cr.

Out of the six metals analyzed only the standard levels for Zn (60 mg/kg), Cu (40 mg/kg) and Cr (2.3 mg/kg) were available for comparison obtained from FAO and WHO permissible levels for food as reported by Charlseet. al, (2011).

The Zn concentrations in the samples were high. *Carica Papaya* [1] recorded highest value of 225 mg/kg while *Colocasia Esculenta* and *Psidium guajava* recorded lowest (75 mg/kg). In general, the samples were all above the standard level of 60 mg/kg. These suggest that Zn is most accumulated metal by plants grown in the area. Zn is an essential element in plants, but could become toxic and pose health issues when consumed in excess.

In the analyzed samples Cu levels ranged from 3.0-4.25 mg kg⁻¹, which is very low when compared to FAO/WHO permissible level for food. However, the concentration of Cu recorded (0.25-3.52) mg kg⁻¹ in the present study is higher than the mean concentration of Cu (2.00 ±1.174) mg kg⁻¹ observed by Aktaruzzaman et al, (2013). Alam et al. (2003) reported mean Cu concentrations of 15.5mg kg⁻¹ and 8.51 mg kg⁻¹ in leafy and non-leafy vegetables respectively from Samata village, Jessor, which is higher than the present findings. Higher mean concentration of Cu (22.19-76.50) mg kg⁻¹ was reported in leafy vegetable species from Turkey (Demirezen and Aksoy, 2006).

The mean levels of Cr in vegetables in the present study were significantly lower than FAO/WHO limit values. It was observed that lower concentrations of Cr were recorded in the present study, when compared with other studies. Gupta et al., (2008) recorded mean concentration of (34.83-96.30 mg kg⁻¹) Cr in West Bengal, India. Sharma et al. (2007) observed mean Cr level of (5.37-27.83 mg kg⁻¹) in Varanasi, India, while Charlse et. al, (2011) recorded values ranging from 1.15-29.29 mg/kg in vegetables grown in the vicinity of the closed dumpsite, which were all higher than the result of the present study. The low concentration of Cr in the area could be due to leaching capabilities of the metal to deeper part of the soil (Banks et. al, 2006).

Table 6. Daily metal intakes estimates for participant in the area

Plant samples	Metals						Mean	SDV
	Cr	Cu	Fe	Mn	Al	Zn		
UL (mg/ day)	N/A	10	8-18	11	N/A	40		
<i>Carica Papaya</i> [1]	0.006	1.2	6.56	0.2	1.68	90	16.61	36.03
<i>ColocasiaEsculenta</i>	0.028	1.42	0.42	-	1.2	30	6.61	13.086
<i>Psidiumguajava</i>	0.034	1.7	1.3	-	-	30	25.36	16.23
<i>Prinusdulcis</i>	0.04	1.6	1.54	0.6	2	40	7.63	12.44
<i>Carica Papaya</i> [2]	0.06	1.68	4.08	0.6	0.42	60	11.14	11.18

High concentration of Fe was recorded in the leaves of edible plants grown in the abandoned. The levels of Fe in plants were significant. However, there was no significant difference in concentration of Fe in plants of the same species. *Carica papaya* [1] recorded the highest Fe level in the study while *Colocasia esculenta* recorded lowest (Table 4). High levels of Fe have been reported in plants grown in abandoned solid waste dumpsites in Port Harcourt, Nigeria (Tanee *et al.*, 2015). The level of Mn recorded in the present study ranged from 0.5 mg/kg in *Carica papaya* [1] to 1.50 mg/kg in both *Prinus dulcis* and *Carica papaya* [2] with mean value of 1.17 mg/kg. Mn was not detected in *Colocasia esculenta* and *Psidium guajava*. In the case of Al, the values recorded are as follows: 1.50 mg/kg in *Carica papaya* [2] to 5 mg/kg in *Prinus dulcis* with mean value of 3.32 mg/kg.

Table 6 is the daily metal intakes estimates for participant in the area. This study estimated the approximate daily metal intake for the participants living in the study area. Experts have recommended a daily intake of at least 0.4 mg of vegetables (Sharma *et al.*, 2008). The daily metals intakes estimate of the metals (Al, Cr, Cu, Fe, Mn and Zn) from each vegetable in this study were calculated as stated in the methodology and result shown in Table 6. The DImetal were compared with the upper tolerable daily intakes for the metals (UL) (Garcia-Rico *et al.*, 2007). In general, Zn recorded high intake estimates for the various metals in the edible plant leave samples. Zinc is required by protein kinases that participate in signal transduction processes and as a stimulator of transducting factors responsible for regulating gene expression (ATSDR, 2005). Zinc plays an important role in the immune system and is an antioxidant *in vivo* (Michael *et al.*, 2009; Strachan, 2010). Deficiency of zinc in human body can result in growth retardation, neuropsychiatric disturbances, dermatitis, alopecia, diarrhoea, increased susceptibility to infections, and loss of appetite (Demirezen and Aksoy, 2006; Michael *et al.*, 2009). Ranking in the descending order of Zn daily intake for the various samples studied; *Carica papaya* [1] > *Carica papaya* [2] > *Prinus dulcis* > *Colocasia esculenta* > *Psidium guajava*.

Cu, Fe and Mn were below the upper tolerable daily intakes for the metals (UL) of 10, 8-18 and 11 mg/day respectively. However, value obtained for Mn in this study were similar to values reported in literature for vegetables by Harmanescu *et al.* (2011) and Uwahet *al.* (2011). Copper levels were very low. Lower copper uptake in human consumption can cause a number of symptoms which include growth retardation, skin ailments, and gastrointestinal disorder (ATSDR, 2004). Copper deficiency also exerts an effect on iodine metabolism resulting in hypothyroidism in animal (Michael *et al.*, 2009).

Table 7. Transfer factor of heavy metal from soils to the leaves of the edible plants

Samples	Transfer factor					
	Cr	Cu	Fe	Mn	Al	Zn
A	7.14×10^{-5}	0.17	0.08	-	42	160.71
B	3.04×10^{-4}	0.22	5.20×10^{-5}	-	37.5	93.75
C	0.88×10^{-4}	0.23	0.03	-	-	-
D	6.67×10^{-4}	0.29	0.03	0.08	31.25	-
E	5.36×10^{-4}	0.34	0.06	0.08	10.5	150
Mean	<0.000	0.25	0.05	0.08	30.31	134.82

Table 7 is the Transfer factor of heavy metal from soils to the leaves of the edible plants. Transfer factor (TF) is the ratio of the concentration of heavy metal in a plant to the concentration of heavy metal in soil. It signifies the amount of heavy metals in the soil that were accumulated in the plant leaves or vegetables (Harrison and Chirgawi, 1989; Smith *et al.*, 1996). This was calculated to understand the extent of risk and hazard associated due to ingestion consequent upon heavy metal accumulation in edible portion of vegetables. Since, soil-to-plant transfer is one of the key components of human exposure to metals through food chain. The transfer factors for each heavy metal were computed based on the method described by Harrison and Chirgawi, (1989). In the present study, the Transfer factor of different heavy metal from soil to vegetable is presented in Table 5. The TF value ranges were: Cr: 7.14×10^{-5} in site A to 6.67×10^{-4} in site D, Cu: 0.17 in site A to 0.34 in site E, Fe: 5.20×10^{-5} in site B to 0.08 in A, Mn: 0.08 in both site D and E respectively, Al: 10.5 in site E to 42 in site A and Zn 93.75 in site B to 160.71 in site A and the trend of TF for heavy metal in vegetable samples studied were in order: Zn>Al>Cu>Mn>Fe>Cr. The physicochemical properties of soil and vegetables could influence the transfer or mobility of metals from soil to plant and is altered by innumerable environmental and human factors (Aktaruzzaman et al, 2013). The highest mean TF value 134.82 and 30.31 was for Zn and Al respectively. However, zinc was more significant than that of aluminium.

4. CONCLUSION

The current status of heavy metal contamination in the abandoned municipal waste dumpsite in Owerri has been investigated. Levels of metals in the soil samples of the study areas were considerably significant. Assessment based on contamination factor, pollution load index and potential ecological risk indexes revealed that the sites are unpolluted with these metals, indicating that the soils were safe for use. Levels of metals in the edible plant leaves are in the order of: Zn>Fe>Cu>Al>Mn>Cr. Zn in particular was higher than FAO/WHO recommended limit. On daily basis, high intake of Zn will be obtained in *Carica Papaya* [1] and [2]. Zn and Al were most transferred from soil to the plant. Therefore, abandoned solid waste dumpsites which contain significant concentrations of heavy metals are later absorbed and accumulated by plants growing on such sites.

References

- [1] Aktaruzzaman, M., A.N.M. Fakhrudin, M.A.Z. Chowdhury, Z. Fardous and M.K. Alam, (2013). Accumulation of Heavy Metals in Soil and their Transfer to Leafy Vegetables in the Region of Dhaka Aricha Highway, Savar, Bangladesh. *Pakistan Journal of Biological Sciences*, 16: 332-338.
- [2] Alam, M.G.M., Snow E.T. and Tanaka, A. (2003). Arsenic and heavy metal contamination of vegetables in Samta village, Bangladesh. *Sci. Total Environ.* 308: 83-96

- [3] Albores, A.F., Perez-Cid, B., Gomes, E.F. and Lopez, E.F. (2000). Comparison between sequential extraction procedures and single extraction procedures and for metal partitioning in sewage sludge samples. *Analyst*, 125, 1353-1357.
- [4] Aluko, O.O. and Oluwande, P.A. (2003). Characterization of leachates from a municipal solid waste landfill site in Ibadan, Nigeria. *Journal of Environmental Health Research*, 2: 83-84
- [5] Awokunmi, E.E., Asaolu S.S. and Ipinmoroti, K.O. (2010). Effect of leaching on heavy metals concentration of soil in some dumpsites. *Afr. J. Environ. Sci. Technol.* 4: 495-499.
- [6] Ayari F, Hamdi H, Jedidi N, Gharbi N and Kossai R (2010). Heavy metal distribution in soil and plant in municipal solid waste compost amended plots. *International Journal Environment, Science and Technology* 7: 465-472.
- [7] Banks MK, Schwab AP, Henderson C. (2006). Leaching and reduction of chromium in soil as affected by soil organic content and plants. *Chemosphere*. 62(2): 255-64.
- [8] Cabrera F, Clemente L, Barrientos DE. (1999). Heavy metal pollution of soils affected by the guadiamar toxic flood. *The Science of the Total Environment*. 242(1-3): 117-129.
- [9] Charles K., William J.S. Mwegoha, Riziki S. Shemdoe (2011). Heavy metals concentrations in vegetables grown in the vicinity of the closed dumpsite. *International Journal Of Environmental Sciences*, 2(2): 889-895.
- [10] Chukwuocha-AC Ngozi. B, Onwuso George. I and Ajoku, Uchechukwu G (2015). Heavy Metals Concentration of Dumpsites and Their Influence on the Soil Physical Properties in Three Major Cities of South Eastern Nigeria. *Journal of Environment and Earth Science*, Pp. 181-196.
- [11] Demirezen, D. and A. Aksoy, (2006). Heavy metal contamination of urban soils and street dusts in limits for Cu, Zn, Ni and exceeded for Cd and Pb. *J. Food Qual.* 29: 252-265.
- [12] Duru C. E., Duru I. A., Ibe F. C. and Enedoh M. C. (2017). Profiling of Zn²⁺ Ion Sorption in Modeled Aqueous Solutions by different Parts of Maize Biomass, *IOSR Journal of Applied Chemistry*, Vol. 10, Issue 3 Ver. 1, 70075.
- [13] Ebong G.A, Akpan M.M, Mkpennie V.N. (2008). Heavy metal contents of municipal and rural dumpsite soils and rate of accumulation by Carica papaya and Talinumtriangulare in Uyo, Nigeria. *E-Journal of chemistry* 5: 281-290.
- [14] Ebong G.A, Etuk H.S, Johnson A.S (2007). Heavy metals accumulation by Talinumtriangulare grown on waste dumpsites in Uyo metropolis, AkwaIbom State, Nigeria. *Journal of Applied Sciences* 7: 1404-1409.
- [15] Elaigwu SE, Ajibola VO, and Folaranmi FM (2007). Studies on the impact of municipal waste dumps on surrounding soil and air quality of two cities in northern Nigeria. *Journal of Applied Sciences* 7: 421-425.
- [16] Enyoh, C.E., Ihionu E.A., Verla A.W., and Ebosie P. N. (2017). Physicochemical Parameter of Palm Oil and Soil from Ihube Community, Okigwe, Imo State Nigeria. *International Letters Of Natural Sciences*, 62, 35-43

- [17] Garcia, W.J., Blessin, C.W., Inglett, G.E. and Kwolek, W.F. (1981). Metal accumulation and crop yield for a variety of edible crops grown in diverse soil media amended with sewage sludge. *Environmental Science Technology*, 15(7), 793-804.
- [18] Garcia-Rico, L., Leyva-Perez, J. & Jara-Marini, M.E. (2007). Content and daily intake of copper, zinc, lead, cadmium, and mercury from dietary supplements in Mexico. *Food and chemical toxicology*, 45(9), 1599-1605.
- [19] Goorah S, Esmiot M, Boojhawon R (2009). The health impact of Non-hazardous solid waste disposal in a community: The case of the Mare Chicose landfill in Mauritius. *Journal of Environmental Health* 72: 48-54.
- [20] Gupta, N., D.K. Khan and S.C. Santra, (2008). An assessment of heavy metal contamination in vegetables grown in wastewater-irrigated areas of Titagarh, West Bengal, India. *Bull. Environ. Cont. Toxicol.* 80: 115-118.
- [21] Hakanson, L., (1980). An ecological risk index for aquatic pollution control. A sedimentological approach. *Water Res.* 14: 975-1001.
- [22] Harmanescu, M., Alda, L. M., Bordean, D.M., Gogoasa, I. and Gergen, I. (2011). Heavy metals health risk assessment for population via consumption of vegetables grown in old mining area; a case study: Banat County, Romania. *Chemistry Central Journal*, 5, 64.
- [23] Harrison, R.M., and Chirgawi, M.B. (1989). The assessment of air and soil as contributors of some trace metals to vegetable plants. Use of a filtered air growth cabinet. *Science of the Total Environment*, 83(1-2), 13-34.
- [24] Jiang X., W. X. Lu, H. Q. Zhao, Q. C. Yang and Z. P. Yang (2014). Potential ecological risk assessment and prediction of soil heavy-metal pollution around coal gangue dump. *Nat. Hazards Earth Syst. Sci.* 14, 1599-1610.
- [25] Kachenko, A.G. and Singh, B. (2006). Heavy metals contamination in vegetables grown in urban and metal smelter contaminated sites in Australia. *Water Air Soil Pollut.* 169: 101-123.
- [26] Leah Amor S. Cortez and Johnny, A. C. (2014). Heavy Metal Concentration of Dumpsite Soil and Accumulation in *Zea mays (corn)* Growing in a Closed Dumpsite in Manila, Philippines. *International Journal of Environmental Science and Development*, Vol. 5, No. 1, 78-80.
- [27] Li, X.D., C.S. Poon and P.S. Liu, (2001). Heavy metal contamination of urban soils and street dusts in Hong Kong. *Applied Geochem.* 16: 1361-1368.
- [28] Ndukwu, B.C., G.C. Obute and E. Eze, (2008). Uptake and accumulation of heavy metals by plants on abandoned refuse dumpsites in parts of Rivers State, Nigeria. *Scientia Africana*, 7: 130-140.
- [29] Njoku, P.C. and Ibe, F.C. (2009). Heavy metal Accumulation in Electronic Technicians, *J. Chem. Soc. Nigeria*, Vol. 34, No 1, 1-4.
- [30] Obasi NA, Akubugwo EI, Ugboogu OC, and Chinyere GC (2012). Heavy Metals Bioavailability and Phyto-accumulation Potentials of Selected Plants on Burrow-pit

- Dumpsites in Aba and Ntigha Dumpsite in IsialaNgwa of Abia State, Nigeria. *Nigerian Journal of Biochemistry and Molecular Biology* 27: 27-45.
- [31] Obute, G.C., Ndukwu, B. C, Eze E (2010). Changes in species diversity and physico-chemical properties of plants in abandoned dumpsites in parts of Por-Harcourt, Nigeria. *Scientia Africana* 9: 181-193.
- [32] Okoronkwo NE, Ano AO and Onwuchekwa EC (2005). Environment, health and risk assessment: a case study of the use of an abandoned municipal waste dump site for agricultural purposes. *African Journal of Biotechnology*, 4 (11): 1217-1221.
- [33] Okoronkwo, N.E., Odemelam, S.A., Ano, O.A (2006). Levels of toxic elements in soils of abandoned waste dumpsite. *African Journal of Biotechnology*, 5(13), pp. 1241-1244.
- [34] Oluyemi EA, Feuyit G, Oyekunle JAO, Ogunfowokan AO (2008). Seasonal variations in heavy metal concentrations in soil and some selected crops at a landfill in Nigeria. *Afr J Environ SciTechno* 12(5): 89-96
- [35] Pekey, H.(2006). The distribution and sources of heavy metals in Izmit Bay surface sediments affected by a polluted stream. *Mar. Pollut. Bull.* 52: 1197-1208.
- [36] Sharma, R.K., M. Agrawal and F. Marshall, (2007). Heavy metal contamination of soil and vegetables in suburban areas of Varansi, India. *Ecotoxicol. Environ. Safety*, 66: 258-266.
- [37] Sharma, R.K., Agrawal, M. and Marshall, F.M. (2008). Atmospheric deposition of heavy metals (Cu, Zn, Cd and Pb) in Varanasi City, India. *Environmental Monitoring and Assessment*, 142, 269-278.
- [38] Shi, G., Z. Chen, S. Xu, J. Zhang, L. Wang, C. Bi and J. Teng, (2008). Potentially toxic metal contamination of urban soils and roadside dust in Shanghai, China. *Environ. Pollut.* 156: 251-260.
- [39] Strachan, S. (2010). Points of view: Nutrition. Trace elements. *Current Anaesthesia and Critical Care*, 21(1), 44-48.
- [40] Tanee F.B.G. and Eshalomi-Mario T.N. (2015). Heavy Metal Contents in Plants and Soils in Abandoned Solid Waste Dumpsites in Port Harcourt, Nigeria. *Research Journal of Environmental Toxicology* 9 (6): 342-349.
- [41] Tomlinson DC, Wilson DJ, Harris CR, Jeffrey DW. Problem in heavy metals in estuaries and the formation of pollution index. *Helgol. Wiss. Meere-sunlter* 1980, 33 (1-4): 566-575.
- [42] Uwah, E. I., Ndahi, N. P., Abdulrahman, F. I. And Ogugbuaja, V. O. (2011). Heavy metal levels in spinach (*Amaranthus caudatus*) and lettuce (*Lactuca sativa*) grown in Maiduguri, Nigeria. *Journal of Environmental Chemistry and Ecotoxicology*, 3(10), 264-271.
- [43] Verla Andrew Wirnkor, Enyoh Christian EbereOkonkwo Felix Chukwunonso, Verla Evelyn Ngozi (2017a). Radiation levels of dumpsites within Imo State University, Owerri, Imo State, Nigeria. *World News of Natural Sciences* 11, 45-64.

- [44] Verla Evelyn Ngozi, Verla Andrew Wirnkorand Enyoh Christian Ebere (2017 b). Pollution assessment models of surface soils in Port Harcourt city, Rivers State, Nigeria. *World News of Natural Sciences* 12, 1-20.
- [45] Wong, C.S.C., Li, X.D., Zhang, G., Qi, S.H., and Peng, X.Z (2003). Atmospheric depositions of heavy metals in the Pearl River Delta, China. *Atmospheric Environment*, 37, pp. 767–776.
- [46] Xiong, Z.T. (1998). Lead uptake and effects on seed germination and plant growth in a lead hyperaccumulator *Brassica pekinensis*. *Rupr Bulletin of Environmental Contamination and Toxicology*, 60, 285-291.
- [47] Tessier, A., Campbell, P. G. C. and Bisson, M. 1979. *Anal. Chem.*, 51: 844–851
- [48] Adams, S. J. and Alloway, B. J. 1988. *Environ. Technol. Letts.*, 9: 695–702
- [49] Hetland, S., Martinsen, I., Radzuk, B. and Thomassen, Y. 1991. *Anal. Sci.*, 7: 1029–1032
- [50] Banat, K., Förstner, U. and Müller, G. 1972. *Naturwissenschaften*, 12: 525–528.
- [51] McLaren, R.G. and Crawford, D.V. 1973. *J. Soil Science*, 24: 172–181.
- [52] Grimme, H. 1967. *Z. Pflanzenernähr. Düng. Bodenk*, 116: 207–222.
- [53] Shuman, L.M. 1979. *Soil Science*, 127: 10–17.
- [54] Goldberg, E.D. and Arrhenius, G.O.S. 1958. *Geochim. Cosmochim. Acta*, 13: 153–212.
- [55] Hirst, D.M. and Nicholls, G.D. 1958. *J. Sediment. Petrol*, 28: 461–468.
- [56] Loring, D.H. and Nota, D.J.G. 1968. *J. Fish. Res. Board Can*, 25: 2327–2347.
- [57] Skei, J. and Paus, P.E. 1979. *Geochim. Commochim. Acta*, 43: 239–246.
- [58] Chester, R. and Hughes, M.J. 1967. *Chem. Geol*, 2: 249–262.
- [59] Agemian, H. and Chau, A.S.Y. 1976. *The Analyst*, 101: 761–767.
- [60] Jenne, E.A., Ball, J.W. and Simpson, C. 1974. *J. Environ. Quality*, 3: 281–287.
- [61] Malo, B.A. 1977. *Environ. Sci. Technol*, 11: 277–282.
- [62] Deurer, R., Förstner, U. and Schmoll, G. 1978. *Geochim. Cosmochim. Acta.*, 42: 425–427
- [63] Presley, B.J., Kolodny, Y., Nissenbaum, A. and Kaplan, I.R. 1972. *Geochim Cosmochim. Acta*, 36: 1073–1099.
- [64] Nissenbaum, A. 1974. *Isr. J. Earth Sci*, 23: 111–116.
- [65] Bruland, K.W., Bertine, K., Koide, M. and Goldberg, E.D. 1974. *Environ. Sci Technol*, 8: 425–432.
- [66] Gibbs, R. 1973. *Science*, 180: 71–73.
- [67] Gibbs, R. 1977. *Geol. Soc. Amer. Bull*, 88: 829–843.
- [68] Gupta, S.K. and Chen, K.Y. 1975. *Environ. Letters*, 10: 129–158.

[69] Fillipek, L.H. and Owen, R.M. 1979. *Chem. Geol.*, 26: 105–117.

[70] Tessier, A., Campbell, P.G.C. and Bisson, M. 1979. *Anal. Chem*, 51: 844–851

(Received 05 July 2017; accepted 24 August 2017)