



# World News of Natural Sciences

An International Scientific Journal

WNOFNS 20 (2018) 121-128

EISSN 2543-5426

---

---

## Effect of Pre-chilling and Storage Temperature on Seed Germination of *Solanum macrocarpon* L. (African Eggplant)

**Damilola Grace Ogunrotimi\*** and **Joshua Kayode**

Department of Plant Science and Biotechnology, Faculty of Science, Ekiti State University, Ado Ekiti, Nigeria

\*E-mail address: [damilolarotimi91@gmail.com](mailto:damilolarotimi91@gmail.com) , +2348064066205

### ABSTRACT

This study examined the effects of storage temperature and pre-chilling on the germination of the seeds of *Solanum macrocarpon* – which, according to literature, have a seed viability of 63.33%. In our study, the seeds were sown in dry Petri dishes and placed in the refrigerator set at 4 °C for 1, 3, 5, 7, and 10 days. Another set of seeds was placed in incubators set at 25, 35, and 45 °C for 40 days before the germination experiment was carried out on a laboratory workbench. Results from the study revealed that while the storage temperature had a significant effect on that the germination of seeds of this plant, pre-chilling of seeds and incubator treatments generally reduced seed germination. Thus, the seed germination in this plant is temperature dependent.

**Keywords:** Germination, Pre-chilling, Seeds, *Solanum macrocarpon*, Temperature

### 1. INTRODUCTION

*Solanum macrocarpon* L., commonly known as African eggplant or Gboma and “Igbagba” among the Yoruba ethnic group of Southwestern Nigeria, is an erect herbaceous perennial herb. It grows up to about 1-1.5 m tall. It is thought to be a native of Africa although it is now widely distributed in different ecological zones. The leaves are broad with pointed apex and arranged in an alternate pattern. The shape of the leaves can be said to be oval and lobed with a wavy margin, both sides of the leaves are hairy with simple hairs.

Prickles are usually found along the mid-rib. The flowers are usually of about 2-3.5 cm and of purple colour, fruits are fleshy, round and flattened at the bottom and contain numerous seeds of reniform shape which are used for propagation (Oboh *et al.*, 2005). The leaves are eaten as a separate dish or sauce together with other ingredients including pepper, tomato, onions, locust beans and fish or meat. It contains vitamins and minerals including vitamins A, C, E, folic acid, niacin, potassium, thiamin, and riboflavin which contribute to growth and the maintenance of good health (Sanstead *et al.*, 1998; Roberts *et al.*, 2000). Appreciable amounts of antioxidants, such as flavonoids, phenolics, ascorbic acid, and tocopherols, which are also found in *S. macrocarpon*, have been shown to be scavengers of harmful radicals that are known to cause cellular damage, heart diseases, cancer, Parkinson's and Alzheimer's diseases (Sodipo *et al.*, 2012). Its leaves are also rich in dietary fiber, an important nutrient found only in plant food.

Seed germination is one of the critical phases in plant life cycle (Shoab *et al.*, 2012) which is affected by dormancy and environmental factors. Adequate temperature, water, light, and moisture content are important environmental factors that are important for seed germination and seedling development. A proper investigation of germination requirements show how a species germination process is adapted to habitat conditions and regulated by environmental factors (Van Assche *et al.*, 2002). It is common for seed lots to have moisture contents considered inadequate for safe and effective storage, such as contents higher than 12%. Reduction in moisture content is important to preserve the physiological quality of seeds for at least eight months; this will slow down possible chemical and physical changes that may come about during storage (Barrozo *et al.*, 2014; Carvalho *et al.*, 2016).

Temperature is considered one of the crucial factors for the process of germination (Forcella, 1998). Before domestication of a plant species can be achieved, proper information about the best environmental condition for its germination must be available. Vegetable crops with orthodox seeds may present differences in germination due to possible dormancy, loss of viability due to preservation conditions, genetic differences among the materials, or aging (Gisbert *et al.*, 2011). Poor seed germination rate limits the use of different species for culture or plant breeding, and different treatments such as scarification, stratification and addition of different chemical substances are commonly used to promote germination in several species (Bone, 2003; Brady and McCourt, 2003). Ecological studies of important plant species such as vegetables provide prerequisite information that will promote their cultivation. Dormancy and low germination rates have been observed in the genus *Solanum* (Adebola and Afolayan, 2006). Recent initiative revealed that very little research has been carried out on seed germination of *Solanum macrocarpon*.

The present study therefore seeks to provide information on the effect of pre-chilling and storage temperature on germination of *Solanum macrocarpon*.

## **2. MATERIALS AND METHODS**

### **Seed collection, weight and moisture content determination**

Mature fruits of *Solanum macrocarpon* were harvested during field surveys from different areas in Ekiti State, Nigeria. The freshly harvested seeds were removed from their capsules, air-dried, and later used for germination studies in this experiment. Seed weight was determined by weighing 100 seeds using an analytical balance and the mean weight of one seed

was calculated. Moisture content of seed was determined by oven drying method and calculated using the formula:

$$\% \text{ Moisture content} = \frac{FW - DW}{FW} \times 100$$

where: FW = Fresh weight of seeds  
DW = Dry weight of seeds.

### Viability testing

Seeds of *S. macrocarpon* to be tested for viability were surface sterilized using 1.0% sodium hypochlorite solution for 5 min and thoroughly rinsed with distilled water. 9 cm sterile Petri-dishes were overlaid with cotton wool and moistened with 7 mL of distilled water. 50 randomly selected seeds were counted, put in the Petri dish and then covered. Treatments were arranged in a complete randomized design and replicated three times. Emergence of radicle was taken as a criterion of seed germination and the germinated seeds were recorded on the 14<sup>th</sup> day. Mean values obtained were separated using the Duncan Multiple Range Test (DMRT) at a probability level of 0.05.

Another batch of seeds were be subjected to floatation method of viability test. This involved steeping of seeds into water in a beaker; seeds that sank to the bottom were identified to be viable.

### Germination experiment

Prior to germination test, seeds were also subjected to refrigerator set at 4 °C for 1, 3, 5, 7, and 10 days. Another set of seeds was placed in incubators set at 25, 35, and 45 °C for 40 days before the germination experiment was conducted on a laboratory work bench at ambient temperatures (AOSA, 1998). Germination tests were conducted in 9-cm sterile Petri dishes lined with cotton wool, moistened with distilled water. Each treatment had three replicates of 20 seeds each. The Petri dishes were examined daily and seeds were considered germinated when their radicles were visible. Germination counts were taken on the 14<sup>th</sup> day after planting. Germination percentage for each treatment was calculated using the formula cited by Czabator (1962):

$$\text{Germination (\%)} = \frac{\text{Total number of seeds germinated}}{\text{Total number of seeds per replicate}} \times 100$$

All experiments were conducted in a complete randomized design with three replicates. Data collected were subjected to analysis of variance with storage temperature and pre-chilling as treatments using SPSS Version 20.

## 3. RESULTS AND DISCUSSION

Seeds of *S. macrocarpon* subjected to viability test yielded 63.33% (**Table 1**). Previous study by Finkeltein *et al.* (2008) revealed the presence of gibberellic acid in freshly extracted

seeds of this species which has been found to enhance germination. Thus the moderately high germination obtained in this study can be attributed to the presence of this acid.

The average seed weight and moisture content of *S. macrocarpon* was  $4.15 \pm 0.09$  mg and  $8.27 \pm 2.70\%$ , respectively (Table 1). Several authors have stated that seed attributes are greatly influenced by climatic and edaphic factors in the addition to density and of plant and genetic variation among species population (Pathak *et al.*, 1975, Rutger and Crowder, 1967). A slight change in seed moisture content greatly affects its storage life (Mane and Puri, 2013). Studies have revealed that a number of orthodox seeds require moisture content of between 6 and 10% for them to retain viability after a long storage (Luna and Wilkinson, 2009). In the present study the moisture content of *Solanum macrocarpon* seeds allows for long term storage without reducing their viability (Table 1).

The effect of pre-chilling and storage temperature on seed germination produced is presented in **Table 2**. The result from the present study shows some variability in germination percentage in both treatments, however, while significant differences were observed in germination of seeds subjected to temperature treatments, no significant statistical differences were obtained in seeds pre-chilled in the refrigerator.

Analysis of variance also shows that pre-chilling has no significant effect on the seed germination of *Solanum macrocarpon* (**Table 3**). A study of the effect of pre-chilling on seed germination of *Solanum nigrum* by Bvenura (2014) revealed a lower germination percentage (28%) in seeds subjected to pre-chilling for 3 days. In the pre-chilling treatment, germination was highest in control seeds followed by seeds pre-chilled for 7 days. Some studies have shown that pre-chilling breaks seed dormancy and enhances germination in plant species (Baskin *et al.*, 2001) while others posit that it has harmful effects on viable seeds (Ren and Tao, 2004).

The cold pre-chilling may change the hormonal balance of seeds and increase germination through enhancement of gibberellic acid and cytokinin activity and/or the decline of abscisic acid (Copeland and McDonald, 2001). Amini *et al.* (2015) also stated that pre-chilling provides enough moisture to activate the hydrolytic enzymes which make seeds ready to germinate once they were moved to the warm temperature. The results from this study however shows that germination reduces with increase in length of days seeds are subjected to pre-chilling, but dropped in seeds that were pre-chilled for 10 days.

In the present study, analysis of variance shows that seed germination is significantly affected by temperature (**Table 4**). The importance of temperature on seed germination has been articulated by some authors, such as Nkomo and Kambizi (2009), Sowunmi and Afolayan (2015).

A variety of high temperature treatments has been employed successfully to overcome dormancy in seeds (Suthar *et al.*, 2009). In the present study, *Solanum macrocarpon* had 10% germination when subjected to a temperature of 25 °C, however Nkomo and Kambizi (2009) observed no germination in *Corchorus olitorius* seeds subjected to a temperature of 25 °C.

Percentage seed germination increased with temperature up to 45 °C, although germination was higher in the control. This is similar to the report of Bvenura (2014) who observed 56% germination in freshly extracted *Solanum nigrum* seeds. Germination was significantly higher in seeds placed in the incubator set at 45 °C. The increase in seed germination percentage which was directly proportional to temperature increase is expected, this agrees with Hegarty (1977) who asserted that the rate of germination increases linearly with temperature within a well defined range.

**Table 1.** Weight, moisture content and viability of *Solanum macrocarpon* seeds

Parameters	Values
Seed weight	4.15 ± 0.09 mg
Seed moisture content	8.27 ± 2.70%
Seed viability	63.33 ± 0.04%

**Table 2.** Effect of pre-chilling and temperature on seed germination of *Solanum macrocarpon*

Treatment	Percentage (%) germination
Pre-chilling 1 day	48.33 ± 25.22 <sup>a</sup>
3 days	33.33 ± 18.78 <sup>a</sup>
5 days	26.67 ± 6.67 <sup>a</sup>
7 days	50.00 ± 7.64 <sup>a</sup>
10 days	25.00 ± 10.00 <sup>a</sup>
Control	61.67 ± 8.82 <sup>a</sup>
Temperature 25 °C	10.00 ± 5.77 <sup>a</sup>
35 °C	18.33 ± 3.33 <sup>a</sup>
45 °C	50.00 ± 5.77 <sup>b</sup>
Control	61.67 ± 8.82 <sup>b</sup>

Mean with different letters along the same column represent significant difference at  $p < 0.05$   
 Values shown are mean ± standard error.

**Table 3.** Analysis of variance of seed germination for pre-chilling treatment in *S. macrocarpon*

Source of variation	SS	Df	MS	F-value	Sig.
Between Group	3245.83	5	649.17	1.02	0.447

Within Group	7616.67	12	634.72		
Total	10862.50	17			

\*Significant at 0.05

**Table 4.** Analysis of variance of seed germination for pre-chilling treatment in *S. macrocarpon*

Source of variation	SS	Df	MS	F-value	Sig.
Between Group	5516.67	3	1838.89	15.76	0.001
Within Group	933.33	8	116.67		
Total	6450.00	11			

\*Significant at 0.05

The degree of enzymatic actions is usually altered by temperature, in addition, Mader (1993) found out that an increase in temperature of 10 °C doubled the rate of enzymatic activity in a living cell. This suggests that *Solanum macrocarpon* seeds will only germinate at optimum temperature suitable for its growth.

#### 4. CONCLUSION

This study suggests that *Solanum macrocarpon* is well able to thrive under varied temperatures, however, cold stratification by pre-chilling is not necessary as the seed does not possess any form of dormancy.

#### References

- [1] Adebola, P.O. and Afolayan, A. J. (2006). Germination responses of *Solanum aculeastrum*, a medicinal species of the Eastern Cape, South Africa. *Seed Science Technology*, 34: 735-740.
- [2] Amini, V., Zaefarian, F., and Rezvani, M. (2014). Effect of pre-chilling and environmental factors on breaking seed dormancy and germination of three foxtail species. *Acta Agriculturae Slovenica*, 105(2): 269-278, DOI: 10.14720/aas.2015.105.2.10
- [3] Barrozo, M.A.S.; Mujumdar, A., and Freire, J.T. (2014) Air-Drying of seeds: A review. *Drying Technology: An International Journal*, 32(10): 1127-1141.

- [4] Bone, M. (2003). Germination of woody legumes from green seed. *Proceedings of International Plant Propagation Society Conference*, 53: 372.
- [5] Brady, S.M. and McCourt, P. (2003). Hormone cross-talk in seed dormancy. *Plant Growth Regulation*, 22: 25-31.
- [6] Bvenura, C. (2014). Wild vegetables of the Eastern Cape of South Africa: The Nutritional Value and Domestication of *Solanum nigrum* L. PhD Thesis, University of Fort Hare, South Africa
- [7] Carvalho, E.R. Oliveira, J.A., Mavaieie, D.P.R., Silva, H.W., and Lopes, C.G.M. (2016) Pre-packing cooling and types of packages in maintaining physiological quality of soybean seeds during storage. *Journal of Seed Science*, 38(2): 129-139. <http://dx.doi.org/10.1590/23171545v38n2158956>.
- [8] Chinedu, S.N., Olasumbo, A.C., Eboji, O.K., Emiloju, O.C., Arinola O.K., and Dania, D.I. (2011). Proximate and Phytochemical Analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. Fruits, *Research Journal of Chemical Sciences*, 1(3): 63-71.
- [9] Czabator, F.J. (1962). Germination value: an index combining speed and completeness of pine seed germination. *Forensic Science*, 8, 386-396.
- [10] Baskin, C.C., Milberg, P., Anderson, L., and Baskin, J.M. (2001). Seed dormancy breaking and germination requirements of *Drosera anglica*, an insectivorous species of the Northern Hemisphere. *Acta Oecologica*, 12: 1-8.
- [11] Copeland, L.O. and McDonald, M., (2001). Principles of seed science and Technology. Kluwer Academic Publisher, Norwell, MA, USA, DOI: 10.1007/9781-4615-1619-4.
- [12] Finkelstein, R., Reeves, W., Ariizumi T., and Steber, C. (2008). Molecular aspects of seed dormancy. *Annual Review of Plant Biology*, 59, 387- 415.
- [13] Forcella, F. (1998). Real-time assessment of seed dormancy and seedling growth for weed management, *Seed Science Research*, 8(2): 201-210.
- [14] Gisbert, C., Prohens, J., and Nuez, F. (2011). Treatments for Improving Seed Germination in Eggplant and Related Species, Proceedings of 5th International Symposium on Seed, Transplant and Stand Establishment of Horticultural Crops Eds.: IA. Pascual and F, Pérez-Alfocea Acta Horticulturae, 898, ISHS
- [15] Hegarty, T.W. (1977). Temperature coefficient (Q<sub>10</sub>), seed germination and other biological processes. *Nature*, 243: 305-306.
- [16] Luna, T. and Wilkinson, K.M. (2009). Collecting, processing and storing seeds. In Dumroese RK, Luna K and Landis TD (Eds). Nursery manual for native plants. a guide for tribal nurseries. Vol I, Nursery Management. Interior, Environment, and Related Agencies Appropriations for 2010: Hearings Before a Subcommittee of the Committee on Appropriations, House of Representatives, One Hundred Eleventh Congress, First Session, Part 3.
- [17] Mader, S.S., 1993. Biology Part 1: The cell (4<sup>TH</sup> Ed) Duburque, IA, USA: Wm. C. Brown communications, Inc. pp. 152.

- [18] Mane, V. and Puri, V. (2013). Studies on permittivity and moisture content of sunflower seeds using microstrip equilateral triangle patch antenna. *International Journal of Engineering Research and Applications*, 3(4): 1584-1586.
- [19] Nkomo, M. and Kambizi, L. (2009). Effects of pre-chilling and temperature on seed germination of *Corchorus olitorius* L. (Tiliaceae) (Jew's Mallow), a wild leafy vegetable, *African Journal of Biotechnology*, 8(6): 1078-1081.
- [20] Pathak, P.S., Dedroy, R., and Rai, P. (1975). Autecology of *Leucaena leucocephala* (L and M) Dewit. J. Seed Polymorphism and germination. *Tropical Ecology*, 15: 1-10.
- [21] Ren, J. and Tao, L. (2004). Effects of different pre-sowing seed treatments on germination of 10 *Calligonum* species. *Forest Ecology and Management*, 195: 291-300
- [22] Roberts, K.M., Daryl, K.G., Peter, A.M., and Victor, W.K. (2000). Harper's Biochemistry, 25th Edition, Lange Medical Book, Appleton and Lange, pp. 209-210.
- [23] Rutger, J.N. and L.V. Crowder, (1967). Effect of plant density on silage and grain yield of six corn hybrids. *Crop. Science*, 7: 182-184.
- [24] Sandstead, H. H., Penland, J.G., Alcock, N.W., Dayal, H.H., Chen, X.C., Li, J.S., Zhao, F., and Yang, J.J. (1998) Effects of Repletion with Zinc and Other Micronutrients on Neuropsychologic Performance and Growth of Chinese Children. *American Journal of Clinical Nutrition*, 68 (2): 470S-475S.
- [25] Shoab, M., Tanveer A., Khaliq A., and Haider Ali H. (2012). Effect of Seed Size and Ecological Factors on Germination of *Emex spinosa*. *World Applied Science Journal*, 17: 964-969.
- [26] Sodipo, O.A. Abdulrahman, F.I. Akan J.C., and Akinniyi, J.A. (2008). Phytochemical Screening and Elemental Constituents of the Fruit of *Solanum macrocarpon* L. *Continental Journal of Applied Sciences*, 3: 85-94.
- [27] Sowunmi, L.I. and Afolayan, A.J. (2015). Effect of environmental factors and sowing depth on seed germination in *Cleome gynandra* L. (Capparaceae). *Pakistan Journal of Botany*, 47(6): 2189-2193.
- [28] Suthar, A. C., Naik, V.R., and Mulani, R.M. (2009). Seed and seed germination in *Solanum nigrum* Linn. *American-Eurasian Journal of Agriculture and Environmental Science*, 5(2): 179-183.
- [29] Van Assche, J., Van Nerum, D., and Darius, P., 2002. The comparative germination ecology of nine *Rumex* species. *Plant Ecology*, 159: 131-142, DOI: 10.1023/A:1015553905110