



# World News of Natural Sciences

An International Scientific Journal

WNOFNS 25 (2019) 220-226

EISSN 2543-5426

---

---

## Seeding cell number required for optimal lipid accumulation during adipocyte differentiation using 3T3-L1 cell line

**Eko Fuji Ariyanto<sup>1,2,\*</sup>, Naren A/L S. Odaiyappan<sup>3</sup>, Lynna Lidyana<sup>4</sup>,  
Tenny Putri Wikayani<sup>5</sup>, Nurul Qomarilla<sup>5</sup>, Dwi Wahyudha Wira<sup>1</sup>,  
Rima Destya Triatin<sup>1</sup>**

<sup>1</sup> Division of Biochemistry and Molecular Biology,  
Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran,  
Jl. Raya Bandung Sumedang, KM.21, Sumedang, Indonesia

<sup>2</sup> Research Center for Medical Genetics, Faculty of Medicine, Universitas Padjadjaran,  
Jl. Eyckman No. 38, Bandung, Indonesia

<sup>3</sup> Faculty of Medicine, Universitas Padjadjaran,  
Jl. Raya Bandung Sumedang, KM.21, Sumedang, Indonesia

<sup>4</sup> Department of Psychiatry, Faculty of Medicine, Universitas Padjadjaran / Dr. Hasan Sadikin General  
Hospital, Jl. Pasteur No.38, Bandung, Indonesia

<sup>5</sup> Cell Culture Laboratory, Faculty of Medicine, Universitas Padjadjaran,  
Jl. Eyckman No. 38, Bandung, Indonesia

\*E-mail address: [fuji@unpad.ac.id](mailto:fuji@unpad.ac.id)

### ABSTRACT

Obesity is one of the major causes of metabolic diseases, such as diabetes and heart attack, and, hence, can lead to low quality of life. Elaborating adipocyte differentiation is very crucial for formulating the treatment and prevention of obesity. The objective of this study is to investigate the seeding cell number required to obtain optimum lipid accumulation during adipocyte differentiation using the 3T3-L1 cell line. Two sets of  $5.48 \times 10^4$  (for Day 0 and Day 8 of differentiation), of  $10.96 \times 10^4$  (for Day 8) and of  $21.92 \times 10^4$  (for Day 8) of 3T3-L1 cells were seeded in each wells of a 12-well plate. Isobuthylmethylxanthine (IBMX), Dexamethasone, and Insulin-containing differentiation cocktails were added into the medium at Day 0 for 48 hours. The medium was changed every two days. Day 0 and Day 8 samples were then stained using Oil Red O and were examined under the microscope to observe the lipid droplets (red-coloured). The lipid droplets were quantified by measuring the

absorbance at wavelength of 550 nm. In the study, seeding the number of  $10.96 \times 10^4$  cells produced very significantly higher lipid accumulation, as compared with seeding the number of  $5.48 \times 10^4$  cells. However, doubling the seeding number into  $21.92 \times 10^4$  cells did not increase the lipid droplets significantly. This study found that the optimum seeding number to obtain the maximum lipid droplets during 3T3-L1 adipocyte differentiation was  $10.96 \times 10^4$  cells.

**Keywords:** 3T3-L1 preadipocytes, differentiation, lipid accumulation, seeding cell number

## 1. INTRODUCTION

The prevalence of obesity is still high worldwide. The World Health Organization (WHO) reported that global obesity has nearly tripled since the year 1975.<sup>4</sup> The prevalence of overweight and obesity in South East Asia is also now increasing, ranging from 8% to 30% in adult males and 8 to 52% in adult females.<sup>1</sup> The prevalence of obesity in adults in Indonesia was 14.6%.<sup>1</sup> The prevalence of obesity in West Java province was 10 to 19 %.<sup>3,4</sup> The region with the highest prevalence of obesity in Indonesia was the eastern region.<sup>2</sup> Adipocyte differentiation or adipogenesis plays important role in obese patients. Adipogenesis is increased in an obese person which leads to their obesity.<sup>14</sup> Adipogenesis can occur due to many reasons but very rarely due to genetical alteration.<sup>9,10</sup> Increased adipocyte differentiation occurs in people who undergo a sedentary lifestyle and people who practice a poor diet consisting of high fat foods.<sup>9</sup>

Since obesity can also be said as excess of lipid accumulation in adipocytes, finding out a way to reduce adipocyte formation might give a chance to formulate treatment and prevention of obesity. For that purpose, 3T3-L1 preadipocyte cell lines, which were originally cloned from 3T3 mouse embryo fibroblasts,<sup>12</sup> were used to study the differentiation of the preadipocyte cells to adipocytes *in-vitro* under chemical stimulation, also known as differentiation cocktail.<sup>3,8</sup>

Although numerous studies, regarding adipocyte differentiation, have been performed worldwide, the protocol may vary in different regions and with different origin of cell lines.<sup>5,13</sup> Optimized protocol is required to get the optimum number of differentiated adipocytes. It is very important because in further studies researchers can take advantage of the optimized protocol to identify substance(s) which has/have inhibition effect on adipocyte differentiation, suggesting potential antiobesity effects.<sup>6</sup>

The purpose of this study is to investigate the suitable optimum seeding cell number required to obtain maximum lipid accumulation during adipocyte differentiation using 3T3-L1 preadipocyte cells at Cell Culture and Cytogenetic Laboratory, Advanced Biomedical Laboratory, Universitas Padjadjaran (UNPAD) located in UNPAD Teaching Hospital, Jalan Eijkman No.38, Bandung, West Java, Indonesia.

## 2. MATERIALS AND METHODS

### 2. 1. Research Design

This is a quantitative analytic study. The purpose of this research is to find the optimum seeding cell number in adipocyte differentiation to develop a protocol for the differentiation of 3T3-L1 preadipocyte cells into adipocytes.

## **2. 2. Cell Differentiation**

In conducting this study, we took advantage of 3T3-L1 preadipocyte cell line. Three different media were used for the process of cell differentiation: MDI (IBMX-dexamethasone-insulin)-containing DMEM-10% FBS-P/S (penicillin-streptomycin) which was called as induction medium, 167 nM Insulin containing DMEM-10% FBS-P/S, and empty DMEM-10% FBS-P/S medium.

The MDI induction medium consists of DMEM-10% FBS-P/S 100 mL, 0.5 mM IBMX, 0.25  $\mu$ M dexamethasone, and 1  $\mu$ g/mL insulin. MDI medium should be stored at a temperature of 4 degree Celsius and should be protected from light. Twelve-well plates were used in all experiments in this study.

On day -5, cell passaging and cell counting should be done (processes explained below) and 2 sets of  $5.48 \times 10^4$ ,  $10.96 \times 10^4$ ,  $21.92 \times 10^4$  cells are seeded in each wells of 12-well plate. DMEM-10% FBS-P/S is added to the wells. On day -2, the medium is changed with the same medium (DMEM-10% FBS-P/S). On day 0, for day 8 samples, medium is replaced with MDI medium. For day 0 samples, the cells are fixed using 4% formaldehyde solution and stored at 4 degree Celsius.

The other set of cells (Day 8 samples) are left to differentiate in the MDI medium for 48 hours. On day 2, the medium of day 8 samples is replaced with DMEM-10% FBS-P/S-167 nM Insulin and left to differentiate for 48 hours. On day 4, the medium of day 8 samples is replaced with DMEM-10% FBS-P/S medium and left to differentiate for 48 hours. On day 6, the medium of day 8 samples is replaced with new DMEM-10% FBS-P/S medium and left to differentiate for 48 hours. On day 8, the cells are now fully differentiated and ready to be stained to determine the percentage of lipid accumulation. The cells (Day 8 samples) as well as Day 0 samples (which was already fixed and stored at 4 degree Celsius) are stained with oil red O (explained below). Then the samples are placed in a microplate reader for the number of lipid cells to be counted the lipid accumulation of Day 0 and Day 8 samples were compared.

## **2. 3. Cell Counting using Improved Neubauer method**

Hemocytometer was used to count the cells.<sup>15</sup> Trypan blue dye was added to the cells to identify live cells.

## **2. 4. Oil Red O Staining**

The cells were fixed with 4% formaldehyde before the staining. After being washed with 60% isopropanol, the cells were stained with Oil Red O solution for 15 minutes. The cells were then rinsed with 60% isopropanol again. Lipid accumulation was stained red.

## **2. 5. Oil Red O Quantification**

Isopropanol was added to the wells to elute the red stain. The eluant was then transferred to another 96 well plate and the absorbance was measured at wavelength of 550 nm. Compare the absorbance value between the wells containing different samples.

## **2. 6. Statistical analysis**













The data were presented in mean $\pm$ SD and analyzed using Microsoft Excel 2016.

### 3. RESULTS

#### 3. 1. Qualitative analysis of lipid accumulation

The macroscopic pictures showed that the well plate in day 0 containing  $5.48 \times 10^4$  number of cells had no lipid accumulation, confirming that they were still preadipocytes. Day 8 samples containing  $10.96 \times 10^4$  and  $21.92 \times 10^4$  number of cells had more abundant lipid accumulation, suggesting mature adipocytes, whereas day 8 sample containing  $5.48 \times 10^4$  number of cells had less lipid accumulation (**Figure 1**).


Lipid accumulation in sample with seeding number of  $10.96 \times 10^4$  cells was very significantly higher as compared to sample with seeding number of  $5.48 \times 10^4$  cells, while the difference between day 8 sample of  $10.96 \times 10^4$  cells and  $21.92 \times 10^4$  cells was not very significant. The microscopic pictures with  $40 \times$  and  $100 \times$  magnification support the results from the macroscopic pictures (Figure 1).

	Day 0		Day 8	
<b>Seeding cell number</b>	$5.48 \times 10^4$	$5.48 \times 10^4$	$10.96 \times 10^4$	$21.92 \times 10^4$
<b>Macroscopic</b>				
<b>Microscopic 40 × magnification</b>				
<b>Microscopic 100 × magnification</b>				

**Figure 1.** Macroscopic and microscopic pictures of cells after oil red O staining. The cells were seeded in three different numbers (as indicated in the figure) on Day -5. MDI stimulation was started on Day 0. Oil red O staining was performed on Day 0 (before MDI stimulation) and eight days after differentiation (Day 8).

### 3. 2. Quantitative analysis of lipid accumulation

Figure 2 showed that Day 8 of sample with seeding number  $10.96 \times 10^4$  cells produced the optimum number of lipid staining intensity at the wavelength of 550 nm and confirms that it had very significantly more lipid accumulation as compared to Day 8 of  $5.48 \times 10^4$  cells seeding number. To double the seeding number from  $10.96 \times 10^4$  cells into  $21.92 \times 10^4$  cells did not increase the lipid accumulation very much (Figure 2).



**Figure 2.** Quantification of lipid droplets. Oil Red O staining was dissolved in 2-propanol and the colour intensity was determined by measuring absorbance at wavelength of 550 nm. \* $p < 0.05$ , \*\* $p < 0.01$ . D0: Day 0, D8: Day 8.

## 4. DISCUSSION

The objective of this study was to investigate the optimum preadipocyte seeding number to acquire maximal lipid accumulation during adipocyte differentiation using 3T3-L1 cells. Our study suggested that seeding cell number of  $10.96 \times 10^4$  produced the optimum lipid accumulation during 3T3-L1 differentiation.

Our study also implied that higher seeding number did not mean that it will result in higher lipid accumulation because a very high seeding cell number can also decrease the differentiation efficiency.<sup>11</sup> This is proved in our study when the difference between day 8

sample containing  $5.48 \times 10^4$  seeding cell numbers and day 8 sample containing  $10.96 \times 10^4$  seeding cell numbers were very significant but the difference between day 8 sample containing  $10.96 \times 10^4$  seeding cell numbers and day 8 sample containing  $21.92 \times 10^4$  seeding cell numbers was less significant. This might be explained by a study conducted by Zebisch *et al.* suggesting that higher seeding cell numbers decreased the differentiation efficiency of the cells.<sup>7</sup>

Protocol of 3T3-L1 differentiation might vary among laboratories. Every cell culture laboratory has their own protocol which includes the seeding number, concentration of differentiation cocktail, etc. This study was conducted to find out the optimum seeding cell number for maximal lipid accumulation. This optimum seeding number of cells can be utilized for further study, for example, for identification of herbal extracts which can inhibit adipocyte differentiation.

## **5. CONCLUSION**

The optimum seeding cell number for maximal lipid accumulation during differentiation of 3T3-L1 preadipocytes into mature adipocytes was  $10.96 \times 10^4$  cells.

### **Acknowledgement**

We thanked Dr. Afiat Berbudi (Universitas Padjadjaran) for the kind gift of 3T3-L1 preadipocytes used in this study and Universitas Padjadjaran for the research fund.

### **Funding Source**

This study was funded by Universitas Padjadjaran Research Grant.

## **References**

- [1] Rachmi C.N., Li M., and Alison Baur L. Overweight and obesity in Indonesia: prevalence and risk factors - a literature review. *Public Health* 2017 Jun, 147: 20-29
- [2] Arundhana A.I., Utami A.P., Dwi Muqni A., and Thalavera M.T. Regional differences in obesity prevalence and associated factors among adults. *Mal J Nutr* Vol. 24(2) (2018) 193-201
- [3] Ruiz-Ojeda F.J., Rupérez A.I., Gomez-Llorente C., Gil A., and Aguilera C.M. Cell Models and Their Application for Studying Adipogenic Differentiation in Relation to Obesity: A Review. *Int J Mol Sci* . 2016 Jun 30, 17(7).
- [4] Suemori H., Yasuchika K., Hasegawa K., Fujioka T., Tsuneyoshi N., and Nakatsuji N. Efficient establishment of human embryonic stem cell lines and long-term maintenance with stable karyotype by enzymatic bulk passage. *Biochem Biophys Res Commun* 2006 Jul; 345(3): 926–32



- [5] Marques E.S., Leite T.H., Azeredo C.M., Cunha D.B., and Verly Júnior E. Effective strategies for prevention, control, and treatment of obesity in primary health care setting for adolescents, adults, and elderly people. *Medicine* (Baltimore) 2018 Jun; 97(22): e10925
- [6] Sung J.-H., Chon J.-W., Lee M.-A., Park J.-K., Woo J.-T., and Park Y.K. The anti-obesity effect of *Lethariella cladonioides* in 3T3-L1 cells and obese mice. *Nutr Res Pract* . 2011 Dec; 5(6): 503–10
- [7] Zebisch K., Voigt V., Wabitsch M., and Brandsch M. Protocol for effective differentiation of 3T3-L1 cells to adipocytes. *Anal Biochem*. 2012 Jun 1, 425(1): 88–90
- [8] Pu Y. and Veiga-Lopez A. PPAR $\gamma$  agonist through the terminal differentiation phase is essential for adipogenic differentiation of fetal ovine preadipocytes. *Cell Mol Biol Lett*. 2017 Dec 23; 22(1): 6
- [9] Hausman G.J., Basu U., Wei S., Hausman D.B., and Dodson M.V. Preadipocyte and Adipose Tissue Differentiation in Meat Animals: Influence of Species and Anatomical Location. *Annu Rev Anim Biosci* 2014 Feb, 2(1): 323–351
- [10] Kaur G. and Dufour J.M. Cell lines: Valuable tools or useless artifacts. *Spermatogenesis* 2012 Jan 1, 2(1): 1–5.
- [11] Chavey C., Lagarrigue S., Annicotte J.S., and Fajas L. (2013) Emerging Roles of Cell Cycle Regulators in Adipocyte Metabolism. In: Bastard JP., Fève B. (eds) *Physiology and Physiopathology of Adipose Tissue*. Springer, Paris
- [12] Suemori H., Yasuchika K., Hasegawa K., Fujioka T., Tsuneyoshi N., and Nakatsuji N. Efficient establishment of human embryonic stem cell lines and long-term maintenance with stable karyotype by enzymatic bulk passage. *Biochem Biophys Res Commun*. 2006 Jul; 345(3): 926–32
- [13] Kawai M., Namba N., Mushiake S., Etani Y., Nishimura R., Makishima M., *et al*. Growth hormone stimulates adipogenesis of 3T3-L1 cells through activation of the Stat5A/5B-PPAR $\gamma$  pathway. *J Mol Endocrinol* 2007 Feb 1, 38(1–2): 19–34
- [14] Cadena-Herrera D., Esparza-De Lara J.E., Ramírez-Ibañez N.D., López-Morales C.A., Pérez N.O., Flores-Ortiz L.F., *et al*. Validation of three viable-cell counting methods: Manual, semi-automated, and automated. *Biotechnol Rep* (Amst). 2015 Apr 18; 7: 9-16. doi: 10.1016/j.btre.2015.04.004
- [15] Hudu S.A., Alshrari A.S., Syahida A., and Sekawi Z. Cell Culture, Technology: Enhancing the Culture of Diagnosing Human Diseases. *J Clin Diagn Res* 2016 Mar; 10(3): DE01-5.