



# World News of Natural Sciences

An International Scientific Journal

WNOFNS 27 (2019) 153-173

EISSN 2543-5426

---

---

## Biocontrol potential of trichoderma and yeast against post harvest fruit fungal diseases: A review

Lamenew Fenta<sup>1</sup>, Habtamu Mekonnen<sup>2</sup>, Tsegaye Gashaw<sup>3</sup>

<sup>1</sup>Department of Biology, Assosa University, Assosa, Ethiopia  
E-mail address: Lamefent21@gmail.com

<sup>2</sup>Department of Biology, Bahir Dar University, Bahir Dar, Ethiopia  
E-mail address: habtie2010@gmail.com

<sup>3</sup>Department of Chemistry, Bahir Dar University, Bahir Dar, Ethiopia  
E-mail address: tsegshchem2004@gmail.com

### ABSTRACT

Crop protection is vital to maintain high productivity and high quality of crops. Over the past years, people used different fungicides, herbicides and good agronomical practices to control fungal diseases and pests to increase productivity. However, extensive use of chemicals in controlling pests and diseases resulted in negative impacts on the environment, producing inferior quality and harming consumer health. In recent times, diverse approaches are being used to manage a variety of pathogens for control of plant diseases. Biological control is the alternative approach for disease management that is eco-friendly and reduces the amount of human contact with harmful chemicals and their residues. A variety of biocontrol agents including fungi and bacteria have been identified; In this regard, yeast and trichoderma species are the most researched microbes in biocontrol research area. But, despite the presence of many reports on biocontrol, practicability of the biocontrols requires effective adoption and a better understanding of the intricate interactions among the pathogen, plants and environment towards sustainable agriculture. To this end, this review attempts to find and compile previous works done on the role of trichoderma and yeast as a biocontrol agent against postharvest fungal pathogens. Moreover, this review analyzes the mechanisms of biocontrol activity, their means of application and future prospects on the use biogents and the challenges that encounter during the commercialization process.

**Keywords:** Biocontrol, Trichoderma, Postharvest, Preharvest, Yeast

## **1. INTRODUCTION**

To date, postharvest loss of fruits and vegetables is a major challenge throughout the world. Plant pathogens are among the most important biotic agents causing serious losses and damages to agricultural products. Hence, plant pathogens need to be controlled to ensure food, feed and fiber production quantitatively and qualitatively (Hasan *et al.*, 2013). Decays of fruits and vegetables caused by fungal pathogens account for significant levels of postharvest losses. It is estimated that about 20–25% of the harvested fruits and vegetables are decayed by pathogens during postharvest handling even in developed countries ((FAO, 2011)). Postharvest diseases of fruits mainly spread during sale, transport and storage (Janisiewicz and Korsten, 2002) and result in reduced food supplies, products of poorer quality, economic hardships for growers and higher prices (Monte, 2001). In addition to quality deterioration and economic losses, fruits infected with fungal pathogens can cause health risk since several fungal genera produce mycotoxins. According to the report of Andersen *et al.* (2006), *Penicillium expansum*, an etiological agent of blue mold in a variety of harvested fruits, produces potential carcinogenic metabolites including citrinin, patulin and chaetoglobosins. Moreover, mycotoxins such as aflatoxins, ochratoxins, alternaria and fumonisin were also found to be produced in fruits and vegetables contaminated with fungal genera such as *Aspergillus*, *Alternaria* and *Fusarium* (Sanzani *et al.*, 2016). Traditionally, postharvest spoilage caused by fungal pathogens is mainly controlled through the use of chemical fungicides either in the field or during storage (Vitoratos *et al.*, 2013). However, the use of many synthetic fungicides in postharvest disease control has become unacceptable due to various reasons including emergence of pathogen resistance to many key fungicides, development of new pathogen biotypes, lack of effective alternative fungicides, increasing levels of fungicide residues in agricultural produce, toxicological problems related to human health and negative environmental impacts (Droby, 2006). As a result of, the global trend is shifting towards the search for safer and eco-friendly alternative approaches to control postharvest diseases and decay in fruits and vegetables (Mari *et al.*, 2012). Among the different biological approaches, use of the microbial antagonists like yeasts, fungi, and bacteria is quite promising and gaining popularity (Droby, 2006).

Biological control is defined as the reduction of inoculum density or disease producing activities of a pathogen or parasite in its active or dormant state, by one or more organisms accomplished naturally or through manipulation of the environment or host or antagonist or by mass introduction of one or more antagonists (Baker and Cook, 1974). Biological control is found to be both environmentally and economically sound. Various biocontrol agents such as fungi and bacteria have been identified for the control of postharvest diseases of many fruits and play an important role in sustainable agriculture and management of plant pathogens (Montesinos, 2003; Sobowale *et al.*, 2008; Montesinos and Bonaterra, 2009; Junaid *et al.*, 2013). It has also been investigated that the potential of antagonistic microorganisms to control decay of fruits and vegetables depends on their ability to colonise fruit surfaces and adapt to various environmental conditions (Sharma *et al.*, 2014). Biological control, using antagonistic microbes to control postharvest disease and decay caused by pathogens is nowadays an emerging and attractive option (Dukare *et al.*, 2011; Liu *et al.*, 2013). The use of antagonist microbes in postharvest disease is advantageous over synthetic fungicides in that antagonist microbes do not produce toxic residues, environmentally friendly, safer application method, easy to deliver and economical to produce (Bonaterra *et al.*, 2012).

In the last one decade so many authors reviewed biocontrol of post harvest fruit pathogens using microbial antagonists in general. However, it is very difficult to address all reports of microbes having antagonistic property at a time as many reports are coming from different corners of the scientific world. Therefore, this review is intended to provide a comprehensive understanding of fungal antagonist-mediated postharvest biocontrol systems, including mechanisms of their biocontrol actions, application and the current perspectives of yeast and trichoderma biocontrol agents in controlling fungal pathogens.

## **2. MAJOR POSTHARVEST PATHOGENS AND THEIR PATHOGENECITY**

Several fungal infections are responsible for postharvest disease and loss of fruits and vegetables. Fungal genera, such as *Alternaria*, *Aspergillus*, *Botrytis*, *Fusarium*, *Geotrichum*, *Gloeosporium*, *Monilinia*, *Colletotrichum*, *Penicillium*, *Mucor* and *Rhizopus* are the major postharvest pathogens responsible for fruits and vegetables diseases (Barkai-Golan, 2001). The symptoms of the disease appear in fungal infected crops during the transportation and storage. Postharvest disease development is affected by several factors and processes, including ripening, harvesting, and mechanical injuries. The diseases development is initiated when fungal pathogens germinate and penetrate the host tissue cuticle through wounds and injuries (Alkan and Fortes, 2015). Fungal pathogens use three main routes to penetrate the host tissue: through wounds, natural openings, such as lenticels, stem ends, and pedicels and by direct breaching of the host cuticle (Emery *et al.*, 2000). Pathogens also enter through the lenticels; pedicel-fruit interphase and sometimes reside endophytically in the stem ends. Moreover, the pathogens can also directly penetrate in the host cuticle throughout the fruit growth period. Several plant fungal pathogens genera like *Alternaria*, *Botrytis*, *Botryosphaeria*, *Colletotrichum*, *Lasioidiplodia*, *Monilinia* and *Phomopsis* reside quiescently at the initial introduction site of unripe fruits (Prusky *et al.*, 2013). As a result, these pathogens remain inactive and invisible during the storage until the fruits ripen. But when the fruit begin to ripe, fungal pathogens grow aggressively and kill the host tissues necrotrophically and take nutrients from the host, leading to decomposition of the tissues and decay initiates. The pathogenic fungi may live dormant either endophytically or hemibiotrophically on fruit tissue till ripening. However, during ripening, the intrinsic disease resistance mechanism protecting the fruits from fungal attack becomes weak or inefficient and then fruits become vulnerable to fungal attacks (Prusky *et al.*, 2013). Therefore, postharvest disease biocontrol becomes vital to prevent post harvest crop loss both in quantity and quality.

## **3. POTENTIAL BIOCONTROL AGENTS AS A POTENTIAL SUBSTITUTE FOR CHEMICAL FUNGICIDES: TRICHODERMA AND YEAST AS PROMISING BIOAGENTS**

Biological control using microbial agents has been reported among several alternatives to be an effective approach, to the use of synthetic chemical fungicides for managing postharvest fruit decay (Droby *et al.*, 2009; Spadaro and Gullino, 2004). Indiscriminate use of chemicals to control the pathogens has caused enhancement of overhead costs, accumulation of toxic chemical residues in food chain and soil pollution leading to loss of soil health.

**Table 1.** Some reports on the use of trichoderma and yeast as biocontrol agent against plant pathogens

<b>Antagonist (yeast &amp; trichoderma)</b>	<b>Pathogen</b>	<b>Host Plant</b>	<b>Citation</b>
<i>Trichoderma harzianum</i>	Anthraco nose ( <i>Colletotrichum musae</i> )	Banana	Devi and Arumugam (2005)
	Anthraco nose ( <i>Colletotrichum gloeosporioides</i> )	Mango	Alvindia (2018):
	Gray mold ( <i>Botrytis cinerea</i> )	Grape	Batta (2007)
	Anthraco nose ( <i>Colletotrichum gloeosporioides</i> )	Mango	Sivakumar <i>et al.</i> (2000)
<i>Trichoderma viride</i>	Stem-end rot	lentil	Sharfuddin <i>et al.</i> (2012)
<i>Trichoderma spp</i>	Root rot	Pine	Dar <i>et al.</i> (2011)
	<i>Sclerotium rolfsii</i>	Mango	Bastakoti <i>et al.</i> (2017)
	Chocolate spot disease ( <i>Botrytis fabae</i> Sard)	Faba Bean	Sahile <i>et al.</i> (2011)
	Anthraco nose	Papaya Fruit	Valenzuela <i>et al.</i> (2015); Admasu <i>et al.</i> (2014)
<i>Trichoderma asperellum</i>	-		Santos-Villalobos (2013)
<i>Trichosporon pullulans</i>	Fruit rots	Mango	Pathak (1997)
	Alternaria rot	Cherry	Qin <i>et al.</i> (2004)
	Gray mold ( <i>Botrytis cinerea</i> )	Cherry	Qin <i>et al.</i> (2004)
Yeast spp (as general)	Anthraco nose	Mango	Montiel <i>et al.</i> (2017)
Yeast	<i>Penicillium italicum</i>	Citrus	Da Cunha <i>et al.</i> (2018)

<i>Saccharomyces cerevisiae</i>	<i>Colletotrichum acutatum</i> (fruit drop)	Citrus	Lopes <i>et al.</i> (2015)
<i>Rhodotorula minuta</i>	Sour rot	Citrus	Ferraz <i>et al.</i> (2016)
<i>Candida azyma</i>			
<i>Aureobasidium pullulans</i>			
<i>Rhodotorula glutinis</i>	Gray mold ( <i>Botrytis cinerea</i> )	Tomato	Liu <i>et al.</i> (2013)
<i>Leucosporidium scottii</i>	<i>P. expansum</i> and <i>B. cinerea</i>	Apple	
<i>Candida oleophila</i> (I-182)	<i>Penicillium digitatum</i>	Grapefruit	Droby <i>et al.</i> (2002)
<i>Candida saitoana</i>	<i>Botrytis cinerea</i>	Apple	El-Ghaouth <i>et al.</i> (2003)
<i>Pichia membranaefaciens</i>	<i>Penicillium expansum</i>	Peach	Yang <i>et al.</i> (2011)
<i>Debaryomyces hansenii</i> , <i>Rhodotorula minuta</i> , <i>Cryptococcus laurentii</i> and <i>Cryptococcus diffluens</i>	Anthraxnose	Mango	Luis <i>et al.</i> (2017)

Apart from this, the chemicals tend to become less efficient due to the development of resistance among the pathogens over time scale. Under these circumstances, the use of various ecofriendly biocontrol agents is increasingly being emphasized as an important component of plant disease management. Moreover, chemical measures may establish imbalance in the microbiological community i.e. unfavorable situation for activity of beneficial organisms (Suryawanshi, 2018). So, direct application of antagonist would be safer method for introducing microorganisms into the soil for biological control of soil borne plant pathogen. Many authors reported that trichoderma and yeasts are promising antagonists against many phytopathogenic diseases of crops (Bastakoti *et al.*, 2017; Liu *et al.*, 2013; Antos-Villalobos *et al.*, 2013; Alvindia, 2018; Marques, 2018).

#### 4. ORIGIN OF MICROBIAL ANTAGONISTS

Most of the microbial antagonists naturally reside on fruit and vegetable surfaces. Many effective antagonists have been isolated and characterized as suitable biocontrol agents for the

control of postharvest pathogens from fruits and vegetables (Janisiewicz *et al.*, 2013). In addition to the fruit surface, phyllosphere is a best source of microbes with antagonistic role (Kalogiannis *et al.*, 2006). Many antagonistic microbes were also isolated and characterized from roots and soil. For instance, the yeast *Rhodotorula glutinis*, obtained from the tomato phyllosphere impedes the growth of *Botrytis cinerea*, a causative agent of grey mold on tomato leaves and fruits (Kalogiannis *et al.*, 2006). The yeast *Kloeckera apiculata* isolated from citrus roots, effectively control the postharvest pathogens *Penicillium italicum* on citrus and grapes (Long *et al.*, 2007). Some effective microbial antagonists have also been investigated in unique natural habitats. For instance, the cold-tolerant yeast *Leucosporidium scottii*, an isolate from the Antarctic soil, was identified as an efficient microbial antagonist to *P. expansum* and *B. cinerea* responsible for blue and grey mold of apples, respectively (Vero *et al.*, 2013). Similarly, the marine yeast *Rhodospiridium paludigenum*, an osmotolerant yeast isolate, inhibits *P. expansum* growth on pear fruits, while *Alternaria alternate* inhibits *P. expansum* growth on Chinese winter jujube (Wang *et al.*, 2011). Marine yeasts typically have greater osmotolerance ability compared to yeasts isolated from the fruit surface and therefore be more suitable candidates for use under conditions with high abiotic stress (Hernandez-Montiel *et al.*, 2010). *Trichoderma* is an asexually reproducing fungal genus usually found in soil. They are known to be strong opportunistic invaders, fast growing, prolific producers of spores and also powerful antibiotic producers even under highly competitive environment for space, nutrients, and light (Montero-Barrientos *et al.*, 2011).

These properties collectively make *Trichoderma* ecologically very dominant and ubiquitous strains able to grow in native prairie, agricultural, marsh, forest, salt and desert soils of all climatic zones. Recently, marine *Trichoderma* isolates were characterized to evaluate their potential use as halotolerant biocontrol agents and found effective against *Rhizoctonia solani* inducing systemic defense responses in plants (Gal-Hemed *et al.*, 2011).

## 5. MECHANISMS OF ACTIONS OF MICROBIAL ANTAGONISTS

Several studies have shown the antifungal potential of many microbial antagonists against postharvest fungal pathogens (Nunes, 2012; Gbadeyan *et al.*, 2016; Wisniewski *et al.*, 2016). Biological control can result from interactions between organisms of different types. In the process of interaction, pathogens are antagonized by the presence and activities of other organisms that they encounter. Competition for nutrients and space, antibiosis through antibiotic production, mycoparasitism, production of cell wall lytic enzymes, and induction of host resistance are major biocontrol mechanisms displayed by antagonists (El-Ghaouth *et al.*, 2004; Sharma *et al.*, 2009; Di Francesco *et al.*, 2016). Recent studies have showed the roles of biofilm formation, quorum sensing, alleviation of host oxidative damage and antifungal volatile compound production in suppressing the activity of postharvest fungal pathogens on fruits (Liu *et al.*, 2013). Often, there is more than one mechanism in successful postharvest biological control and some of the different mechanisms of antagonism are mentioned below.

### 5. 1. Competition

Biological agents compete with plant pathogens for space, organic nutrients and minerals. An effective competition for nutrients such as carbohydrates, amino acids, vitamins and minerals as well as for oxygen and/ or space is vital to suppress postharvest pathogens of fruits

(Spadaro *et al.*, 2016). As the major postharvest diseases are caused by fungi, the majority of antagonists are highly efficient biocontrol agents by successfully competing with fungi for nutritional resources (Zhang *et al.*, 2010). Under nutrient starvation, the antagonists diminish the available nutrients in the wound site and make nutrients inaccessible for the pathogens to germinate, grow and infect. The capability of antagonistic yeasts to attach with their pathogen hyphae increases nutrient competition and thus obstructs the initiation process of the pathogenic infection (Talibi *et al.*, 2014). Yeasts can use most of the carbohydrate and nitrogen sources for cell growth by competing for space and forming an extracellular polysaccharide matrix at the wound site and as a result of this, the growth rate of yeast antagonists is generally high (Spadaro *et al.*, 2010). The analysis of the radio-labelled glucose distribution pattern among the antagonistic yeast *Sporobolomyces roseus* and an etiological agent of grey mold disease (*B. cinerea*) revealed strong sugar use by the antagonistic yeast, which ultimately blocked the conidial germination of the pathogen due to sugar deficiency (Spadaro and Gullino, 2004). Similarly, the important role of the competition for sugars and nitrates was observed in the interactions of *Pichia guilliermondii* with *B. cinerea* on apples (Spadaro and Droby, 2016) and *Colletotrichum* spp. on peppers (Chanchaichaovivat *et al.*, 2008). In fruit wounds, competition is extended to other essential nutrients, such as oxygen, amino acids or vitamins when present at low concentrations. A significant decrease in the efficacy of a yeast strain of *A. pullulans*, an antagonist against *P. expansum* was observed when high concentrations of amino acids were applied exogenously to apple wounds (Bencheqroun *et al.*, 2007).

This shows the important role of competition for nutrients, which represents major mechanism in biocontrol activity. Surface residing non-pathogenic natural microbiota of fruits can also intervene in nutrient and space competition by effective colonisation and toxic metabolite production (Galvez *et al.*, 2010; Di Francesco *et al.*, 2016). Further, the rapid colonisation of wound site also depends on the antagonist concentration and the host fruit species, as certain antagonists prefer certain nutrient types. Most aerobic and facultative anaerobic micro-organisms respond to low iron stress by producing extracellular, low molecular weight (500-1000 daltons) iron transport agents, designated as Siderophores, which selectively make complex with iron (Fe<sup>+</sup>) with very high affinity. Iron starvation prevents the germination of spores of fungal pathogens in rhizosphere as well as rhizoplane. Eg. Siderophores produced by *Pseudomonas fluorescens* (known as pseudobactins or pyoverdins) helps in the control of soft rot bacterium, *Erwinia caratovora*. suppression of *Pythium ultimum* by *Enterobacter cloacae*, *P. putida* colonize the root system in the rhizosphere and a corresponding reduction in Fusarium wilt suppression in cucumber (Maindad *et al.*, 2014).

## 5. 2. Induction of host resistance

Plants actively respond to a variety of environmental stimuli, including gravity, light, temperature, physical stress, water and nutrient availability. Several studies have investigated that the application of microbial biocontrol agents to fruit surfaces induced systemic resistance against invading fungal pathogens (Janisiewicz *et al.*, 2008; Romanazzi *et al.*, 2016; Droby *et al.*, 2016). Induction of resistance to biotic or abiotic stresses involves accumulation of structural barriers and elicitation of many biochemical and molecular defense responses in the host, including mitogen activated protein kinase signaling (MAPK), reactive oxygen species generation (ROS), biosynthesis of terpenoid and phytoalexin, production of phytoalexins and PR-proteins, enhanced accumulation of phenolic compounds, lignification at the infection site

and strengthening of host cell wall by formation of glycoproteins, lignin, callose, and other phenolic polymers (Shoresh *et al.*, 2010).

*Trichoderma* spp. are well-known for their ability to promote plant growth and defense. The previous studies showed that *Trichoderma* can produce gluconic and citric acids that decrease the soil pH, enhance the solubilization of phosphates, micronutrients, and mineral components such as iron, magnesium, and manganese (Vinale *et al.*, 2016). It was shown that the bean plants treated with *T. harzianum* T019 always had an increased size respect to control. In addition, this strain induced the expression of plant defense-related genes and produced a higher level of ergosterol, indicating its positive effects on plant growth and defense in the presence of the pathogen (Mayo *et al.*, 2015).

Moreover, the roots of maize plants treated with *T. harzianum* strain T-22 were about twice as long compared to untreated plants after several months from treatment (Harman, 2004). Saravanakumar *et al.* (2016) showed that *Trichoderma cellulase* complexes trigger the induced systematic resistance (ISR) against *Curvularia* leaf spot in maize by increasing the expression of genes related to the jasmonate/ethylene signaling pathways. Furthermore, Rao *et al.* (2015) suggested that treatment of legume seeds (*Cajanus cajan*, *Vigna radiate* and *Vigna mungo*) with *T. viride* induces systemic resistance by reprogramming defense mechanisms in these legumes.

Treatment of peaches with yeast *C. laurentii* and methyl jasmonic acid (MeJA) stimulated the activities of enzymes chitinase, b-1, 3-glucanase, phenylalanine ammonia-lyase (PAL) and peroxidase (POD) in comparison to the application of yeast or MeJA alone (Yao and Tian, 2005). The treatment reduced the diameter of disease lesions on fruit caused by *M. fructicola* and *P. expansum*. Higher levels of the enzymes, such as catalase (CAT), glutathione peroxidase, methionine sulfoxide reductase peroxiredoxin, and polyphenol oxidase (PPO), protect the host tissues against oxidative damage by *P. expansum* pathogen. Application of antagonistic yeast *R. paludigenum* on mandarins at pre-harvest stage also induced defense response by increasing production of defense-related enzymes, including b-1, 3-glucanase (Spadaro *et al.*, 2016).

### 5. 3. Antibiosis

Antagonism mediated by specific or nonspecific metabolites of microbial origin, by lytic agents, enzymes, volatile compounds or other toxic substances is known as antibiosis. Most microbes produce and secrete one or more compounds with antibiotic activity. In some instances, antibiotics produced by microorganisms have been shown to be particularly effective at suppressing plant pathogens and the diseases they cause. *Trichoderma* can produce low molecular weight diffusible compounds or antibiotics that inhibit the growth of other microorganisms. There are several metabolites or antibiotics secreted from *Trichoderma* against their pathogens such as: harzianic acid, tricholin, peptaibols, 6-pentyl- $\alpha$ -pyrone, viridin, gliovirin, glisoprenins, and heptelidic acid (Gajera *et al.*, 2013).

Sadykova *et al.* (2015) tested the antibiotic activity in 42 strains of 8 species of the *Trichoderma* genus (*Trichoderma asperellum*, *T. viride*, *T. hamatum*, *T. koningii*, *T. atroviride*, *T. harzianum*, *T. Citrinoviride*, and *T. longibrachiatum*) isolated from Siberian. Vinale *et al.* (2014) also showed that the pyrone 6-pentyl-2H-pyran-2-one is a metabolite purified from the culture filtrate of different *Trichoderma* spp. (*T. viride*, *T. atroviride*, *T. harzianum* and *T. koningii*) and has shown both *in vivo* and *in vitro* antifungal activities towards several plant pathogenic fungi. Furthermore, Howell (1999) reported that strains of *Trichoderma virens* (P group) produce the antibiotic gliovirin which is very active against *P. ultimum*, while the Q group of these strains can produce gliotoxin, which is very active against *R. solani*.



#### 5. 4. Mycoparasitism

Direct parasitism, mycoparasitism or hyperparasitism, is the ability of antagonistic microorganism to attach with the hyphae of fungal pathogens to produce extracellular cell wall lytic enzymes. Mycoparasitism of antagonist depends upon the sequential occurrence of the following events: come into close contact of fungal pathogens, mutual recognition by antagonist and pathogen, lytic enzymes secretion and, active growth of antagonist into the host (Spadaro and Gullino, 2004; Talibi *et al.*, 2014). Parasitism causes either complete killing of fungal propagules or destruction and lysis of their structure. Wisniewski *et al.* (1991) reported mycoparasitism initially in the studies on biocontrol of *Botrytis cinera* by yeast antagonist *P. guilliermondii*. Disintegration of pathogenic fungal cell wall by the action of extracellular hydrolytic enzymes of antagonists, such as chitinases, chitosanases, glucanases, cellulase and/or protease, individually or in combination, contributes to biocontrol activity (Spadaro and Droby, 2016). The lytic enzymes also impede pathogen spores germination, elongate of germ-tube and destroy oospores (El-Tarabily, 2006). Banani *et al.* (2015) also reported chitinase activity of antagonistic yeast *Metschnikowia fructicola* and demonstrated that chitinase gene *MfChi* was over induced in the presence of yeast *Monilinia fructicola* cell wall. An overexpressed *MfChi* chitinase in *Pichia pastoris* controlled the growth of *M. fructicola* and *Monilinia laxa* under in vitro and in vivo studies on peach fruits. Similarly, antifungal activity of alkaline serine protease, secreted by yeast-like fungus *A. pullulans*, is documented as mycoparasitism (Zhang *et al.*, 2012). Enzymatic breakdown of fungal pathogens hyphae results in cellular deformities, including cytological damages, lysis and distortion in mycelia, altered cell membrane permeability and leakage of cytoplasmic content (Di Francesco *et al.*, 2016). It may, therefore, be inferred that enzymatic dissolution of cell walls leads to the loss of fungal protoplasm and accountable for antagonistic activity (Kim and Chung, 2004).


#### 5. 5. Production of antifungal volatile compounds (VOCs)

Microbial antagonists produce several antifungal metabolites of high VOCs also play important role in inhibition of fungal pathogen growth (Mari *et al.*, 2016). VOCs are low molecular weight lipophilic compounds mixture. Role of VOCs produced by fungi (Morath *et al.*, 2012); yeast (Di Francesco *et al.*, 2015); and bacteria (Zhang *et al.*, 2013) have been reported to control postharvest disease of fruits. Fungal species, such as *Trichoderma harzianum*, *Fusarium oxysporum* and *A. pullulans*, produce volatile antifungal substances in low concentrations (Mari *et al.*, 2012). Yeast antagonist *A. pullulans* produces VOCs, including 2-methyl-1-butanol, 3-methyl-1-butanol, phenethyl alcohol and 2-methyl-1-propanol, which showed fungistatic activity against *C. acutatum*, *B. cinerea* and *Penicillium* species (Di Francesco *et al.*, 2014). Fungi having ability to produce VOCs may be used as bio-fumigant. Endophytic fungus *Muscodora albus*, isolated from *Cinnamomum zeylanicum* in a botanical garden of Honduras, is a typical example of volatile producing bio-fumigant fungi for control of postharvest decay (Strobel, 2011).

### 6. APPLICATION METHODS OF BIOCONTROL AGENTS

Microbial antagonists are used either those that exist on the produce itself which can be promoted and managed or those that can be artificially introduced against post harvest

pathogens (Sharma *et al.*, 2009) (**Figure 1**). Many investigators have provided strong evidence that several pathogens infest fruits and vegetables in the field, and these infestations become critical factors for decays during transportation or storage of the commodities and hence argued that preharvest application (s) of microbial antagonistic culture are often effective to controlling postharvest decays in fruits and vegetables (Ippolito *et al.*, 2004; Irtwange, 2006).



**Fig. 1.** Interaction within the plants, pathogenic microorganisms and biocontrol

Typically, pre-harvest application is done to pre-colonize the fruit surface with the antagonistic microbes so that fruits can be colonized by the antagonists before colonization by the pathogens (Ippolito and Nigro, 2000).

Preharvest applications with yeast antagonists *Tricho sporonpullulans* (Lindner), *Cryptococcus laurentii* (Kuffer.) *Rhodotorula glutinis* (Fresenius) (Tian *et al.*, 2004),

*Trichoderma harzianum* (Zhang *et al.*, 2007) and *Epicoccum nigrum* (Larena *et al.*, 2005) have proved postharvest control of strawberries where synthetic fungicides proved ineffective. In a different study, the preharvest application of *Aureobasidium pullulans* reduced significantly the storage rots in strawberry (Lima *et al.*, 2003), grapes and cherries (Scheda *et al.*, 2003), and apples (Leibinger *et al.*, 1997). There was a related reducing trend of incidence of green mold (*Penicillium digitatum*) on grape fruit by preharvest spray of *Pichia guilliermondii* (Droby *et al.*, 1992). In pear, field application(s) of *Cryptococcus laurentii* and *Candida oleophila* was reported to reduce storage rots (Benbow and Sugar, 1999) whereas preharvest applications of *Pantoea agglomerans* CPA-2 and *Epicoccum nigrum* were reported to be effective against citrus rots and peach brown rot under laboratory and field conditions respectively (Sharma *et al.*, 2009). Similarly, Canamas *et al.* (2008) reported that preharvest application of different concentrations of *Pantoea agglomerans* was effective against *Penicillium digitatum* during storage of oranges.

Postharvest application of antagonistic microorganisms is common and appears to be better for controlling post harvest diseases of fruits and vegetables. In the application process, the antagonists are sprayed directly onto the surfaces of the fruits and vegetables or are applied by dipping (Sharma *et al.*, 2009). Investigations by many authors show that the postharvest application of microbial antagonists for controlling diseases in fruits and vegetables are more effective than the preharvest approach for citrus (Long *et al.*, 2007), apples ( Morales *et al.*, 2008; Zhang *et al.*, 2009; Mikani *et al.*, 2008), peach (Mandal *et al.*, 2007), banana (Lassois *et al.*, 2008), mango (Kefialew and Ayalew, 2008), tomato (Zhao *et al.*, 2008), and cabbage (Adeline and Sijam, 1999). In strawberries and lemons for example, Pratella and Mari (1993) found that postharvest application of *Trichoderma harzianum*, *Trichoderma viride*, *Gliocladium roseum* and *Paecilomyces variotii* Bainier resulted in a better control of *Botrytis* and *Alternaria* rots respectively than preharvest application(s). In a related development, postharvest applications of *Pseudomonas variotii* and *Trichoderma harzianum* were more effective in controlling *Aspergillus* and *Fusarium* rots in lemons and potatoes than their respective dips in iprodion and benomyl. Identification and selection of promising antagonists are generally followed by the selection of the appropriate time and application method for the effective suppression of postharvest pathogens. In general, both pre-harvest and postharvest application approaches are practiced.

## **7. CHALLENGES AND FUTURE PROSPECTS OF COMMERCIAL BIOCONTROL AGENTS**

In the present crop production scenario, the biocontrol is of utmost importance, but its potential is yet to be exploited fully mainly because the research in this area is still confined to the laboratory and very little attention has been paid to produce the commercial formulations of bio agents. Moreover, whatever has been commercially produced has not been used efficiently by the farmers owing to the lack of information regarding its use. In order to be successful, a biocontrol product needs to provide an acceptable and consistent level of control of target diseases in the target commodity under commercial processing and storage conditions. These conditions may vary significantly for different commodities and in different packinghouses (Droby *et al.*, 2009). Pilot studies, semicommercial, and large-scale commercial tests requiring large amounts of formulated product are needed to obtain the data necessary to

evaluate the efficacy of a biocontrol agent (Abadias *et al.*, 2003). It is important for these tests to be conducted in packinghouses at different locations under conditions of natural infection (Long *et al.*, 2007). Regulatory approval of a formulation by government agencies is also required to produce a commercial product. The application package for approval must contain a third-party evaluation of the safety of the formulated product to human health, as well as efficacy data.

The registration of biocontrol products for postharvest use in the USA is the responsibility of the Environmental Protection Agency (EPA) and on average has required about two years. In contrast, the registration process in Europe takes almost seven years (Nunes, 2012). Although many different yeasts, isolated from a variety of sources, have been reported as good postharvest biocontrol agents, only a few yeast-based biocontrol products are available in the market: Shemer™ (*M. fructicola*, Bayer, Germany), Candifruit™ (*C. sake*, Sipcam-Inagra, Spain), and Boni-Protect™ (the yeast-like fungus, *A. pullulans*, Bioprotect, Germany). These products are registered for use on several different commodities and for several different pathogens. The ability to control different rots on different commodities is essential for the economic viability of a postharvest biocontrol product. For example, Shemer, based on a heat-, oxidative stress- and osmo-tolerant strain of *M. fructicola* NRRL Y-27328 (Droby *et al.*, 2009; Kurtzman and Droby, 2001; Liu *et al.*, 2011), has been shown to be effective against rots caused by *Botrytis*, *Penicillium*, *Rhizopus* and *Aspergillus* on strawberry (Karabulut *et al.*, 2004), grape, sweet potato, carrot and citrus (Blachinsky *et al.*, 2007).

Most of the bio-agents perform well in the laboratory conditions but fail to perform to their fullest once applied to the soil. This is probably attributed to the physiological and ecological constraints that limit the efficacy of bio agents. To overcome this problem, genetic engineering and other molecular tools offer a new possibility for improving the selection and characterisation of bio control agents. Various methods that can contribute to increase the efficacy of bio agent include mutation or protoplasm fusion utilising poly ethylene glycol. There is also an urgent need to mass produce the bio agents, understand their mechanism of action and to evaluate the environmental factors that favour the rapid growth of bio control agents.

The first fungus *Trichoderma harzianum* ATCC 20476 was registered with the EPA for the control of plant diseases. Currently, a total of 14 bacteria and 12 fungi have been registered with the EPA for the control of plant diseases (Fravel, 2005). Most of these are sold commercially as one or more products. The technology of commercialization is still in its initial phase. 65% of the EPA registered organisms have been registered within the past 10 years while the remaining 36% registered over the past 5 years. Although the number of bio control products in plant disease management is increasing, these products still represent only 1% of the agricultural control measures while fungicides account for 15% of total chemicals used in agriculture (Fravel *et al.*, 2005).

In the past five decades, an increasing number of chemical fertilizer and biocidal molecules were the main cause for a substantial increase in crop production and quality. Because of environmental issues and health concerns, continuous and extensive use of those molecules has raised serious debate, and often various biological control methods based on natural pest and pathogen-suppressing organisms are being recommended as a substitute. Globally the registrations of microbial biocontrol agents are increasing significantly. The changes in legislation in the country level, development of new policies and management structures to address the reduction of chemical uses are the expanding scope of biocontrol agents.

On the other hand, the researchers worldwide have been supported to discover new biocontrol agents to reinforce for entering in the industry. Being practical, at present biocontrol agents are not comparable to chemical pesticides in meeting efficacy which is needed for market expectations, but they still have a promising future if knowledge and methods of various fields of biotechnology are utilized. The availability of recent molecular technologies has significantly facilitated for surveying and identification of candidate agents, and helped to interpret the modes of action after field applications. These new technologies like proteomics and functional genomics will give new possibilities for insights in ecological constraints and will help to see unseen possibilities to determine the physiological status and expression of crucial genes present within the biocontrol agents during mass production, formulation, storage and application.

## **8. CONCLUSIONS**

Pesticide residues in fresh fruits and vegetables have been, and will continue to be, one of the main concerns of regulatory agencies and consumers. Therefore, reducing or eliminating the pre- and postharvest use of synthetic chemical fungicides by developing alternative management strategies remains a high research priority. This review article has provided a brief overview on the use of yeast and trichoderma antagonists as a viable alternative to synthetic, chemical fungicides. It is anticipated that the continuing withdrawal of key postharvest fungicides from the market, due to exclusion by regulatory agencies or the high-cost of re-registration, will lead to an absence of effective tools for reducing postharvest losses. Hence, the use of biocontrol products is expected to gain momentum in the coming years and become more widely accepted as a component of an integrated approach to managing postharvest diseases. Many challenges need to be addressed in order to develop a commercially successful postharvest biocontrol product. These include the improvement and enhancement of biocontrol efficacy under commercial condition, the development of high quality, economical methods of fermentation and formulation, the maintenance of cell viability and biocontrol efficacy in the formulated product, the identification of yeast and trichoderma antagonists that exhibit a wide spectrum of activity against several different pathogens on different commodities; e) the establishment of an effective marketing outlet, preferably by a multinational based company; and f) developing a fundamental understanding of how biocontrol systems operate and how the environment affects the interactions between the host, pathogen, and biocontrol agent.

## **References**

- [1] Hasan, R. N., Ali, M. R., Shakier, S. M., Khudhair, A. M., Hussin, M. S., Kadum, Y. A., and Abbas, A. A. (2013). Antibacterial activity of aqueous and alcoholic extracts of *Capsella Bursa* against selected pathogenic bacteria. *American Journal of BioScience*, 1(1), 6-10.
- [2] FAO, I., IMF, O., and UNCTAD, W. (2011). The World Bank, the WTO, IFPRI and the UN HLTF (2011). Price Volatility in Food and Agricultural Markets: Policy Responses. Rome, FAO.

- [3] Janisiewicz, W.J. and Korsten, L. (2002). Biological control of postharvest diseases of fruits. *Annual review of phytopathology*, 40(1), 411-441.
- [4] Monte, E. (2001). Understanding Trichoderma: between biotechnology and microbial ecology. *International Microbiology*, 4(1), 1-4.
- [5] Andersen, B. and Thrane, U. (2006). Food-borne fungi in fruit and cereals and their production of mycotoxins. In *Advances in Food Mycology* (pp. 137-152). Springer, Boston, MA.
- [6] Sanzani, S. M., Reverberi, M., and Geisen, R. (2016). Mycotoxins in harvested fruits and vegetables: Insights in producing fungi, biological role, conducive conditions, and tools to manage postharvest contamination. *Postharvest Biology and Technology*, 122, 95-105.
- [7] Vitoratos, A., Bilalis, D., Karkanis, A., and Efthimiadou, A. (2013). Antifungal activity of plant essential oils against *Botrytis cinerea*, *Penicillium italicum* and *Penicillium digitatum*. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 41(1), 86-92.
- [8] Droby, S. (2006). Biological control of postharvest diseases of fruits and vegetables: difficulties and challenges. *Phytopathologia Polonica*, 39, 105-117.
- [9] Mari, M., Martini, C., Spadoni, A., Rouissi, W., and Bertolini, P. (2012). Biocontrol of apple postharvest decay by *Aureobasidium pullulans*. *Postharvest Biology and Technology*, 73, 56-62.
- [10] Baker, K.F. and Cook, R.J. (1974). Biological control of plant pathogens. WH Freeman and Company.
- [11] Montesinos, E. (2003). Development, registration and commercialization of microbial pesticides for plant protection. *International Microbiology*, 6(4), 245-252.
- [12] Sobowale, A.O. and Oyewole, O.B. (2008). Effect of lactic acid fermentation of cassava on functional and sensory characteristics of fufu flour. *Journal of food processing and preservation*, 32(4), 560-570.
- [13] Montesinos, E. and Bonaterra, A. (2009). Microbial pesticides. *Encyclopedia of microbiology*, 3rd edn. Elsevier, New York, 110-120.
- [14] Junaid, M., Adnan, M., Khan, N., Khan, N., and Ali, N. (2013). Plant Growth, Biochemical Characteristics And Heavy Metals Contents Of *Medicago Sativa* L., *Brassica Juncea* (L.) Czern. And *Cicer Arietinum* L. *Fuuast Journal of Biology*, 3(2), 95-103.
- [15] Sharma, S. D., Kumar, P., and Yadav, S. K. (2014). Glomus–Azotobacter association affects phenology of mango seedlings under reduced soil nutrient supply. *Scientia horticulturae*, 173, 86-91.
- [16] Dukare, A. S., Prasanna, R., Dubey, S. C., Nain, L., Chaudhary, V., Singh, R., and Saxena, A. K. (2011). Evaluating novel microbe amended composts as biocontrol agents in tomato. *Crop protection*, 30(4), 436-442.

- [17] Liu, P., Luo, L., and Long, C. A. (2013). Characterization of competition for nutrients in the biocontrol of *Penicillium italicum* by *Kloeckera apiculata*. *Biological control*, 67(2), 157-162.
- [18] Bonaterra, A., Badosa, E., Cabrefiga, J., Francés, J., and Montesinos, E. (2012). Prospects and limitations of microbial pesticides for control of bacterial and fungal pomefruit tree diseases. *Trees*, 26(1), 215-226.
- [19] Droby, S., Wisniewski, M., Macarisin, D., and Wilson, C. (2009). Twenty years of postharvest biocontrol research: is it time for a new paradigm? *Postharvest biology and technology*, 52(2), 137-145.
- [20] Barkai-Golan, R. (2001). Postharvest diseases of fruits and vegetables: development and control. Copyright © 2001 Elsevier. ISBN 978-0-444-50584-2
- [21] Alkan, N. and Fortes, A. M. (2015). Insights into molecular and metabolic events associated with fruit response to post-harvest fungal pathogens. *Frontiers in plant science*, 6, 889.
- [22] Prusky, D., Alkan, N., Mengiste, T., and Fluhr, R. (2013). Quiescent and necrotrophic lifestyle choice during postharvest disease development. *Annual Review of Phytopathology*, 51, 155-176.
- [23] Spadaro, D. and Gullino, M. L. (2004). State of the art and future prospects of the biological control of postharvest fruit diseases. *International journal of food microbiology*, 91(2), 185-194.
- [24] Suryawanshi, K. T. (2018). *Development Of Disease Management Strategies For Minimal Fungicides Residue In Table Grapes* (Doctoral dissertation, Indira Gandhi Krishi Vishwavidhyalaya, Raipur).
- [25] Bastakoti, S., Belbase, S., Manandhar, S., and Arjyal, C. (2017). Trichoderma species as Biocontrol Agent against Soil Borne Fungal Pathogens. *Nepal Journal of Biotechnology*, 5(1), 39-45.
- [26] de los Santos-Villalobos, S., Guzmán-Ortiz, D. A., Gómez-Lim, M. A., Délano-Frier, J. P., de-Folter, S., Sánchez-García, P., and Peña-Cabriales, J. J. (2013). Potential use of *Trichoderma asperellum* (Samuels, Liechfeldt et Nirenberg) T8a as a biological control agent against anthracnose in mango (*Mangifera indica* L.). *Biological Control*, 64(1), 37-44.
- [27] Alvindia, D. G. (2018). The antagonistic action of *Trichoderma harzianum* strain DGA01 against anthracnose-causing pathogen in mango cv. 'Carabao'. *Biocontrol science and technology*, 28(6), 591-602.
- [28] Marques, E., Martins, I., and Mello, S. C. M. D. (2018). Antifungal potential of crude extracts of *Trichoderma* spp. *Biota Neotropica*, 18(1). e20170418. <http://dx.doi.org/10.1590/1676-0611-BN-2017-0418>
- [29] Devi, A. N. and Arumugam, T. (2005). Studies on the shelf life and quality of Rasthali banana as affected by postharvest treatments. *Orissa J. Hortic*, 33(2), 3-6.

- [30] Batta, Y. A. (2007). Control of postharvest diseases of fruit with an invert emulsion formulation of *Trichoderma harzianum* Rifai. *Postharvest Biology and technology*, 43(1), 143-150.
- [31] Sivakumar, D., Wijeratnam, R. W., Wijesundera, R. L. C., Marikar, F. M. T., and Abeyesekere, M. (2000). Antagonistic effect of *Trichoderma harzianum* on postharvest pathogens of rambutan (*Nephelium lappaceum*). *Phytoparasitica*, 28(3), 240.
- [32] Sharfuddin, R. Y. and Mohanka, N. A. (2012). In vitro antagonism of indigenous *Trichoderma* isolates against phytopathogen causing wilt of lentil. *International Journal of Life Science & Pharma Research*, 2, 195-202.
- [33] Dar, G. H., Beig, M. A., Ahanger, F. A., Ganai, N. A., and Ahangar, M. A. (2011). Management of root rot caused by *Rhizoctonia solani* and *Fusarium oxysporum* in blue pine (*Pinus wallichiana*) through use of fungal antagonists. *Asian J. Plant Pathol*, 5(2), 62-67.
- [34] Sahile, S., Sakhuja, P. K., Fininsa, C., and Ahmed, S. (2011). Potential antagonistic fungal species from Ethiopia for biological control of chocolate spot disease of faba bean. *African Crop Science Journal*, 19(3), 213-225.
- [35] Valenzuela, N. L., Angel, D. N., Ortiz, D. T., Rosas, R. A., García, C. F. O., and Santos, M. O. (2015). Biological control of anthracnose by postharvest application of *Trichoderma* spp. on maradol papaya fruit. *Biological Control*, 91, 88-93.
- [36] Admasu, W., Sahile, S., and Kibret, M. (2014). Assessment of potential antagonists for anthracnose (*Colletotrichum gloeosporioides*) disease of mango (*Mangifera indica* L.) in North Western Ethiopia (Pawe). *Archives of Phytopathology and Plant Protection*, 47(18), 2176-2186.
- [37] Pathak, V. N. (1997). Mundkur Memorial Lecture-Post-harvest Fruit Pathology-- Present Status and Future Possibilities. *Indian Phytopathology*, 50(2), 161-185.
- [38] Qin, G., Tian, S., and Xu, Y. (2004). Biocontrol of postharvest diseases on sweet cherries by four antagonistic yeasts in different storage conditions. *Postharvest biology and technology*, 31(1), 51-58.
- [39] Hernandez Montiel, L. G., Zulueta Rodriguez, R., Angulo, C., Rueda Puente, E. O., Quiñonez Aguilar, E. E., and Galicia, R. (2017). Marine yeasts and bacteria as biological control agents against anthracnose on mango. *Journal of Phytopathology*, 165(11-12), 833-840.
- [40] da Cunha, T., Ferraz, L. P., Wehr, P. P., and Kupper, K. C. (2018). Antifungal activity and action mechanisms of yeasts isolates from citrus against *Penicillium italicum*. *International journal of food microbiology*, 276, 20-27.
- [41] Lopes, M. R., Klein, M. N., Ferraz, L. P., da Silva, A. C., and Kupper, K. C. (2015). *Saccharomyces cerevisiae*: a novel and efficient biological control agent for *Colletotrichum acutatum* during pre-harvest. *Microbiological research*, 175, 93-99.
- [42] Ferraz, L. P., da Cunha, T., da Silva, A. C., and Kupper, K. C. (2016). Biocontrol ability and putative mode of action of yeasts against *Geotrichum citri-aurantii* in citrus fruit. *Microbiological research*, 188, 72-79.



- [43] Liu, J., Sui, Y., Wisniewski, M., Droby, S., and Liu, Y. (2013). Utilization of antagonistic yeasts to manage postharvest fungal diseases of fruit. *International journal of food microbiology*, 167(2), 153-160.
- [44] Droby, S., Vinokur, V., Weiss, B., Cohen, L., Daus, A., Goldschmidt, E. E., and Porat, R. (2002). Induction of resistance to *Penicillium digitatum* in grapefruit by the yeast biocontrol agent *Candida oleophila*. *Phytopathology*, 92(4), 393-399.
- [45] El Ghaouth, A., Wilson, C. L., and Wisniewski, M. (2003). Control of postharvest decay of apple fruit with *Candida saitoana* and induction of defense responses. *Phytopathology*, 93(3), 344-348.
- [46] Yang, Z., Cao, S., Cai, Y., and Zheng, Y. (2011). Combination of salicylic acid and ultrasound to control postharvest blue mold caused by *Penicillium expansum* in peach fruit. *Innovative Food Science & Emerging Technologies*, 12(3), 310-314.
- [47] Janisiewicz, W. J., Jurick II, W. M., Vico, I., Peter, K. A., and Buyer, J. S. (2013). Culturable bacteria from plum fruit surfaces and their potential for controlling brown rot after harvest. *Postharvest biology and technology*, 76, 145-151.
- [48] Kalogiannis, S., Tjamos, S. E., Stergiou, A., Antoniou, P. P., Ziogas, B. N., and Tjamos, E. C. (2006). Selection and evaluation of phyllosphere yeasts as biocontrol agents against grey mould of tomato. *European journal of plant pathology*, 116(1), 69-76.
- [49] Long, C. A., Deng, B. X., and Deng, X. X. (2007). Commercial testing of *Kloeckera apiculata*, isolate 34-9, for biological control of postharvest diseases of citrus fruit. *Annals of Microbiology*, 57(2), 203.
- [50] Vero, S., Garmendia, G., González, M. B., Bentancur, O., and Wisniewski, M. (2013). Evaluation of yeasts obtained from Antarctic soil samples as biocontrol agents for the management of postharvest diseases of apple (*Malus domestica*). *FEMS yeast research*, 13(2), 189-199.
- [51] Wang, Y., Tang, F., Xia, J., Yu, T., Wang, J., Azhati, R., and Zheng, X. D. (2011). A combination of marine yeast and food additive enhances preventive effects on postharvest decay of jujubes (*Zizyphus jujuba*). *Food chemistry*, 125(3), 835-840.
- [52] Montero-Barrientos, M., Hermosa, R., Cardoza, R. E., Gutiérrez, S., and Monte, E. (2011). Functional analysis of the *Trichoderma harzianum* nox1 gene, encoding an NADPH oxidase, relates production of reactive oxygen species to specific biocontrol activity against *Pythium ultimum*. *Appl. Environ. Microbiol.* 77(9), 3009-3016.
- [53] Gal-Hemed, I., Atanasova, L., Komon-Zelazowska, M., Druzhinina, I. S., Viterbo, A., and Yarden, O. (2011). Marine isolates of *Trichoderma* spp. as potential halotolerant agents of biological control for arid-zone agriculture. *Appl. Environ. Microbiol.* 77(15), 5100-5109.
- [54] Nunes, C. A. (2012). Biological control of postharvest diseases of fruit. *European Journal of Plant Pathology*, 133(1), 181-196.
- [55] Gbadeyan, F. A., Orole, O. O., and Gerard, G. (2016). Study of naturally sourced bacteria with antifungal activities. *Int. J. Microbiol. Mycol* 4(1), 9-16.

- [56] Wisniewski, M., Droby, S., Norelli, J., Liu, J., and Schena, L. (2016). Alternative management technologies for postharvest disease control: the journey from simplicity to complexity. *Postharvest Biology and Technology*, 122, 3-10.
- [57] Di Francesco, A., Martini, C., and Mari, M. (2016). Biological control of postharvest diseases by microbial antagonists: how many mechanisms of action? *European Journal of Plant Pathology*, 145(4), 711-717.
- [58] Spadaro, D. and Droby, S. (2016). Development of biocontrol products for postharvest diseases of fruit: the importance of elucidating the mechanisms of action of yeast antagonists. *Trends in Food Science & Technology*, 47, 39-49.
- [59] Zhang, J. and Zhou, J. M. (2010). Plant immunity triggered by microbial molecular signatures. *Molecular Plant*, 3(5), 783-793.
- [60] Zhang, D., Spadaro, D., Garibaldi, A., and Gullino, M. L. (2010). Efficacy of the antagonist *Aureobasidium pullulans* PL5 against postharvest pathogens of peach, apple and plum and its modes of action. *Biological Control*, 54(3), 172-180.
- [61] Talibi, I., Boubaker, H., Boudyach, E. H., and Ait Ben Aoumar, A. (2014). Alternative methods for the control of postharvest citrus diseases. *Journal of applied microbiology*, 117(1), 1-17.
- [62] Spadaro, D. and Gullino, M. L. (2004). State of the art and future prospects of the biological control of postharvest fruit diseases. *International journal of food microbiology*, 91(2), 185-194.
- [63] Spadaro, D. and Droby, S. (2016). Development of biocontrol products for postharvest diseases of fruit: the importance of elucidating the mechanisms of action of yeast antagonists. *Trends in Food Science & Technology*, 47, 39-49.
- [64] Chanchaichaovivat, A., Panijpan, B., and Ruenwongsa, P. (2008). Yeast biocontrol of a fungal plant disease: a model for studying organism interrelationships. *Journal of biological education*, 43(1), 36-40.
- [65] Bencheqroun, S. K., Bajji, M., Massart, S., Labhilili, M., El Jaafari, S., and Jijakli, M. H. (2007). In vitro and in situ study of postharvest apple blue mold biocontrol by *Aureobasidium pullulans*: evidence for the involvement of competition for nutrients. *Postharvest Biology and Technology*, 46(2), 128-135.
- [66] Gálvez, A., Abriouel, H., Benomar, N., and Lucas, R. (2010). Microbial antagonists to food-borne pathogens and biocontrol. *Current opinion in biotechnology*, 21(2), 142-148.
- [67] Romanazzi, G., Sanzani, S. M., Bi, Y., Tian, S., Martínez, P. G., and Alkan, N. (2016). Induced resistance to control postharvest decay of fruit and vegetables. *Postharvest Biology and Technology*, 122, 82-94.
- [68] Shoreh, M., Harman, G. E., and Mastouri, F. (2010). Induced systemic resistance and plant responses to fungal biocontrol agents. *Annual review of phytopathology*, 48, 21-43.
- [69] Vinale, F., Strakowska, J., Mazzei, P., Piccolo, A., Marra, R., Lombardi, N., and Lorito, M. (2016). Cremenolide, a new antifungal, 10-member lactone from *Trichoderma*

- cremeum with plant growth promotion activity. *Natural product research*, 30(22), 2575-2581.
- [70] Mayo, S., Gutierrez, S., Malmierca, M. G., Lorenzana, A., Campelo, M. P., Hermosa, R., and Casquero, P. A. (2015). Influence of *Rhizoctonia solani* and *Trichoderma* spp. in growth of bean (*Phaseolus vulgaris* L.) and in the induction of plant defense-related genes. *Frontiers in Plant Science*, 6, 685.
- [71] Harman, G. E., Howell, C. R., Viterbo, A., Chet, I., and Lorito, M. (2004). *Trichoderma* species—opportunistic, avirulent plant symbionts. *Nature reviews microbiology*, 2(1), 43.
- [72] Saravanakumar, K., Yu, C., Dou, K., Wang, M., Li, Y., & Chen, J. (2016). Synergistic effect of *Trichoderma*-derived antifungal metabolites and cell wall degrading enzymes on enhanced biocontrol of *Fusarium oxysporum* f. sp. *cucumerinum*. *Biological Control*, 94, 37-46.
- [73] Rao, H. Y., Rakshith, D., and Satish, S. (2015). Antimicrobial properties of endophytic actinomycetes isolated from *Combretum latifolium* Blume, a medicinal shrub from Western Ghats of India. *Frontiers in biology*, 10(6), 528-536.
- [74] Yao, H. and Tian, S. (2005). Effects of pre-and post-harvest application of salicylic acid or methyl jasmonate on inducing disease resistance of sweet cherry fruit in storage. *Postharvest Biology and Technology*, 35(3), 253-262.
- [75] Gajera, H., Domadiya, R., Patel, S., Kapopara, M., and Golakiya, B. (2013). Molecular mechanism of *Trichoderma* as bio-control agents against phytopathogen system—a review. *Curr Res Microbiol Biotechnol*, 1(4), 133-142.
- [76] Sadykova, V. S., Kurakov, A. V., Kuvarina, A. E., and Rogozhin, E. A. (2015). Antimicrobial activity of fungi strains of *Trichoderma* from Middle Siberia. *Applied biochemistry and microbiology*, 51(3), 355-361.
- [77] Wisniewski, M., Biles, C., Droby, S., McLaughlin, R., Wilson, C., and Chalutz, E. (1991). Mode of action of the postharvest biocontrol yeast, *Pichia guilliermondii*. I. Characterization of attachment to *Botrytis cinerea*. *Physiological and Molecular Plant Pathology*, 39(4), 245-258.
- [78] El-Tarabily, K. A. and Sivasithamparam, K. (2006). Potential of yeasts as biocontrol agents of soil-borne fungal plant pathogens and as plant growth promoters. *Mycoscience*, 47(1), 25-35.
- [79] Banani, H., Spadaro, D., Zhang, D., Matic, S., Garibaldi, A., and Gullino, M. L. (2015). Postharvest application of a novel chitinase cloned from *Metschnikowia fructicola* and overexpressed in *Pichia pastoris* to control brown rot of peaches. *International journal of food microbiology*, 199, 54-61.
- [80] Zhang, D., Spadaro, D., Valente, S., Garibaldi, A., and Gullino, M. L. (2012). Cloning, characterization, expression and antifungal activity of an alkaline serine protease of *Aureobasidium pullulans* PL5 involved in the biological control of postharvest pathogens. *International journal of food microbiology*, 153(3), 453-464.

- [81] Kim, P. and Chung, K. C. (2004). Production of an antifungal protein for control of *Colletotrichum lagenarium* by *Bacillus amyloliquefaciens* MET0908. *FEMS Microbiology Letters*, 234(1), 177-183.
- [82] Mari, M., Bautista-Baños, S., and Sivakumar, D. (2016). Decay control in the postharvest system: Role of microbial and plant volatile organic compounds. *Postharvest Biology and Technology*, 122, 70-81.
- [83] Morath, S. U., Hung, R., and Bennett, J. W. (2012). Fungal volatile organic compounds: a review with emphasis on their biotechnological potential. *Fungal Biology Reviews*, 26(2-3), 73-83.
- [84] Zhang, X., Li, B., Wang, Y., Guo, Q., Lu, X., Li, S., and Ma, P. (2013). Lipopeptides, a novel protein, and volatile compounds contribute to the antifungal activity of the biocontrol agent *Bacillus atrophaeus* CAB-1. *Applied microbiology and biotechnology*, 97(21), 9525-9534.
- [85] Strobel, G., Singh, S. K., Riyaz-Ul-Hassan, S., Mitchell, A. M., Geary, B., and Sears, J. (2011). An endophytic/pathogenic *Phoma* sp. from creosote bush producing biologically active volatile compounds having fuel potential. *FEMS microbiology letters*, 320(2), 87-94.
- [86] Abadias, M., Usall, J., Teixidó, N., and Viñas, I. (2003). Liquid formulation of the postharvest biocontrol agent *Candida sake* CPA-1 in isotonic solutions. *Phytopathology*, 93(4), 436-442.
- [87] Karabulut, O. A., Romanazzi, G., Smilanick, J. L., and Lichter, A. (2005). Postharvest ethanol and potassium sorbate treatments of table grapes to control gray mold. *Postharvest biology and technology*, 37(2), 129-134.
- [88] Blachinsky, D., Antonov, J., Bercovitz, A., El-ad, B., Feldman, K., Husid, A., and Keren-Zur, M. (2007). Commercial applications of shemer for the control of pre-and post-harvest diseases. *IOBC WPRS BULLETIN*, 30(6/2), 75.
- [89] Fravel, D. R. (2005). Commercialization and implementation of biocontrol. *Annu. Rev. Phytopathol.* 43, 337-359.
- [90] Pratella, G. C., Mari, M., Guizzardi, M., and Folchi, A. (1993). Preliminary studies on the efficiency of endophytes in the biological control of the postharvest pathogens *Monilinia laxa* and *Rhizopus stolonifer* in stone fruit. *Postharvest Biology and Technology*, 3(4), 361-368.
- [91] Adeline, T. S. Y., and Sijam, K. (1999). Biological control of bacterial soft rot of cabbage. In *Biological Control in the Tropics: Towards Efficient Biodiversity and Bioresource Management for Effective Biological Control: Proceedings of the Symposium on Biological Control in the Tropics*. CABI Publishing, Wallingford, UK (pp. 133-134).
- [92] Kefialew, Y., and Ayalew, A. (2008). Postharvest biological control of anthracnose (*Colletotrichum gloeosporioides*) on mango (*Mangifera indica*). *Postharvest Biology and Technology*, 50(1), 8-11.

- [93] Lassois, L., de Bellaire, L. D. L., and Jijakli, M. H. (2008). Biological control of crown rot of bananas with *Pichia anomala* strain K and *Candida oleophila* strain O. *Biological Control*, 45(3), 410-418.
- [94] Mikani, A., Etebarian, H. R., Sholberg, P. L., O’Gorman, D. T., Stokes, S., and Alizadeh, A. (2008). Biological control of apple gray mold caused by *Botrytis mali* with *Pseudomonas fluorescens* strains. *Postharvest Biology and Technology*, 48(1), 107-112.
- [95] Morales, H., Sanchis, V., Usall, J., Ramos, A. J., and Marín, S. (2008). Effect of biocontrol agents *Candida sake* and *Pantoea agglomerans* on *Penicillium expansum* growth and patulin accumulation in apples. *International journal of food microbiology*, 122(1-2), 61-67.
- [96] Cañamás, T. P., Viñas, I., Usall, J., Torres, R., Anguera, M., and Teixidó, N. (2008). Control of postharvest diseases on citrus fruit by preharvest applications of biocontrol agent *Pantoea agglomerans* CPA-2: Part II. Effectiveness of different cell formulations. *Postharvest Biology and Technology*, 49(1), 96-106.
- [97] Benbow, J. M. and Sugar, D. (1999). Fruit surface colonization and biological control of postharvest diseases of pear by preharvest yeast applications. *Plant Disease*, 83(9), 839-844.
- [98] Leibinger, W., Breuker, B., Hahn, M., and Mendgen, K. (1997). Control of postharvest pathogens and colonization of the apple surface by antagonistic microorganisms in the field. *Phytopathology*, 87(11), 1103-1110.
- [99] Schena, L., Nigro, F., Pentimone, I., Ligorio, A., and Ippolito, A. (2003). Control of postharvest rots of sweet cherries and table grapes with endophytic isolates of *Aureobasidium pullulans*. *Postharvest Biology and Technology*, 30(3), 209-220.
- [100] Lima, G., De Curtis, F., Castoria, R., and De Cicco, V. (2003). Integrated control of apple postharvest pathogens and survival of biocontrol yeasts in semi-commercial conditions. *European Journal of Plant Pathology*, 109(4), 341-349.
- [101] Zhang, H., Wang, L., Dong, Y., Jiang, S., Cao, J., and Meng, R. (2007). Postharvest biological control of gray mold decay of strawberry with *Rhodotorula glutinis*. *Biological Control*, 40(2), 287-292.
- [102] Tian, S., Qin, G., and Xu, Y. (2004). Survival of antagonistic yeasts under field conditions and their biocontrol ability against postharvest diseases of sweet cherry. *Postharvest Biology and Technology*, 33(3), 327-331.
- [103] Nunes, C. A. (2012). Biological control of postharvest diseases of fruit. *European Journal of Plant Pathology*, 133(1), 181-196.
- [104] Ippolito, A., Nigro, F., and Schena, L. (2004). Control of postharvest diseases of fresh fruits and vegetables by preharvest application of antagonistic microorganisms. *Crop management and postharvest handling of horticultural products*, 4, 1-30.
- [105] S. V. Irtwange. (2004). Application of Biological Control Agents in Pre- and Postharvest Operations. *Agricultural Engineering International: the CIGR Ejournal. Invited Overview No. 3. Vol. VIII.*